

Distinct degrees and their distribution in complex networks

This content has been downloaded from IOPscience. Please scroll down to see the full text.

J. Stat. Mech. (2013) P06002

(<http://iopscience.iop.org/1742-5468/2013/06/P06002>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 155.41.111.161

This content was downloaded on 26/09/2013 at 15:27

Please note that [terms and conditions apply](#).

Distinct degrees and their distribution in complex networks

P L Krapivsky¹ and S Redner²

¹ Department of Physics, Boston University, Boston, MA 02215, USA

² Center for Polymer Studies and Department of Physics, Boston University, Boston, MA 02215, USA

E-mail: pkrapivsky@gmail.com and redner@buphy.bu.edu

Received 9 April 2013

Accepted 3 May 2013

Published 6 June 2013

Online at stacks.iop.org/JSTAT/2013/P06002

[doi:10.1088/1742-5468/2013/06/P06002](https://doi.org/10.1088/1742-5468/2013/06/P06002)

Abstract. We investigate a variety of statistical properties associated with the number of distinct degrees that exist in a typical network for various classes of networks. For a single realization of a network with N nodes that is drawn from an ensemble in which the number of nodes of degree k has an algebraic tail, $N_k \sim N/k^\nu$ for $k \gg 1$, the number of distinct degrees grows as $N^{1/\nu}$. Such an algebraic growth is also observed in scientific citation data. We also determine the N dependence of statistical quantities associated with the sparse, large- k range of the degree distribution, such as the location of the first hole (where $N_k = 0$), the last doublet (two consecutive occupied degrees), triplet, dimer ($N_k = 2$), trimer, etc.

Keywords: growth processes, network dynamics, random graphs, networks

ArXiv ePrint: [1304.1951](https://arxiv.org/abs/1304.1951)

Contents

1. Introduction	2
2. Distinct degrees	4
3. The first hole	7
4. Last doublet and last dimer	8
5. Discussion	9
Acknowledgments	11
References	11

1. Introduction

A complete microscopic representation of a macroscopic system is usually unavailable and often unnecessary, especially if the system is evolving or it is taken from an ensemble and the goal is to understand the typical features of the ensemble. Thus instead of determining a huge number of parameters (such as the 10^{23} coordinates and momenta of atoms), it often suffices to know a few useful macroscopic quantities (like the total number of atoms and the total energy) to understand the bulk properties of a macroscopic system.

In the realm of networks, one usually starts with an ensemble of large networks that are generated according to a specified and not completely deterministic algorithm. In analogy with other bulk systems, we are typically interested in macroscopic-like network characteristics, such as the total number of links, the total number of triangles, the total number of clusters (maximal connected components), etc [1]. Two of the most useful macroscopic characteristics are the cluster-size distribution and the degree distribution.

The degree of a node (the number of links attached to the node) is perhaps the simplest local network characteristic. It has been now been extensively studied, with an emphasis on networks with broadly distributed degrees [2]. Here we analyze the number of *distinct* degrees D_N that exists for a given network of size N . The number D_N varies from realization to realization, but for the ensembles that we study D_N turns out to be a self-averaging quantity, so that its mean value is the most important characteristic. We focus on $\langle D_N \rangle$ which we generally write as D_N when no ambiguity is possible.

We also investigate the locations of the first hole (the smallest k where N_k equals zero), the last doublet (the largest k value for which $N_k > 0$ and $N_{k+1} > 0$), the last triplet, the last dimer (the largest k value where $N_k = 2$), trimer, etc in the degree distribution (figure 1).

The number of distinct degrees D_N exhibits interesting behavior for network ensembles in which the degree distribution has an algebraic tail; hence we focus on such networks. For concreteness, we consider networks that are grown by preferential attachment. The best-known case is strictly linear preferential attachment [3]–[8], in which a new node attaches to a pre-existing node of degree k with rate $A_k = k$. To illustrate the quantities




Figure 2. The number of nodes of degree $k \geq 10$ for a single network realization of $N = 10^7$ nodes that is grown by strictly linear preferential attachment. The largest degree is $k_{\max} = 6693$, $D_N = 465$, the last doublet occurs at $k = 782$, the last dimer at $k = 641$, the last triplet at $k = 518$, the last trimer at $k = 500$, and the first hole at $k = 201$ (arrows).

provides complementary information about which degree sequences are actually realized in a complex network.

2. Distinct degrees

Consider networks whose degree distribution has the asymptotic power-law form of equation (2). We deal only with sparse networks, for which $\nu > 2$. A network with such a degree distribution can be easily constructed by the redirection algorithm [18], in which a new node either attaches to a randomly selected ‘target’ node with probability $1 - r$ or to the ancestor of the target with probability r . This algorithm generates a scale-free network whose growth rule is precisely shifted linear preferential attachment, with the attachment rate to a node of degree k , $A_k = k + \lambda$, and with $\lambda = 1/r - 2$. This growth rule leads to a degree distribution that has the form (2) with exponent $\nu = 1 + 1/r$. We use this redirection algorithm for our simulations and interchangeably refer to the growth mechanism as either shifted linear preferential attachment or redirection.

To determine the number of distinct degrees that appear in a typical realization of a large network, first notice that for k in the range $k \leq K = (NR)^{1/\nu}$, $N_k \geq 1$. In this dense regime of the degree distribution (figure 2), all degrees with $k < K$ are present. This range therefore gives a contribution of $(NR)^{1/\nu}$ to D_N . In the complementary sparse range of $k > K$, we estimate the number of distinct degrees, by integrating the degree distribution for $k > K$. Adding the contributions from the dense and sparse regimes gives

$$D_N^{\text{naive}} = \frac{\nu}{\nu - 1} K, \quad K = (NR)^{1/\nu}. \quad (3)$$




Figure 3. The average number of distinct degrees D_N versus N for networks that are grown by redirection with redirection probability r . The upper curve (\circ) corresponds to $A_k = k - \frac{1}{2}$ or to redirection probability $r = \frac{2}{3}$. Here the degree distribution exponent is $5/2$ and $D_N = BN^{2/5}$, with $B = (3/2)^{2/5}\pi^{-1/5}\Gamma(3/5) = 1.393019\dots$. The lower curve (\triangle) corresponds to $A_k = k$ or $r = \frac{1}{2}$. Here D_N is given by (5). Each data point represents an average over 10^4 realizations. The dashed lines correspond to the theoretical prediction (1).

While the N -dependence is correct, $D_N \sim N^{1/\nu}$, the amplitude is wrong. A better estimate can be obtained by assuming that the probability distribution for the number of nodes of each degree k is the Poisson distribution with average value N_k given by (2). Then $P_k \equiv \text{Prob}[(\# \text{ nodes of degree } k) \geq 1] = 1 - \exp(-N_k)$. Using this property leads to a more accurate estimate (this same approach was developed in [13])

$$D_N = \sum_{k \geq 1} [1 - e^{-N_k}]. \quad (4)$$

Replacing the sum by an integral, we ultimately obtain (1). For strictly linear preferential attachment, $R = 4$ and $\nu = 3$, so that

$$D_N = BN^{1/3} \quad B = 2^{2/3}\Gamma\left(\frac{2}{3}\right) = 2.149528\dots \quad (5)$$

In contrast, the naive estimate (3) for the amplitude is $B_{\text{naive}} = 3 \times 2^{-1/3} = 2.381101\dots$, which exceeds the more accurate value by $\approx 11\%$.

Generally $D_N/D_N^{\text{naive}} = \Gamma(2 - 1/\nu)$, so for the admissible range of $2 < \nu < \infty$, this ratio monotonically increases from $\frac{1}{2}\sqrt{\pi} \approx 0.886227$ to 1. As shown in figure 3, simulation results are in excellent agreement with our theoretical predictions. A more detailed asymptotic analysis (employing methods developed in [19]) indicates that the average number of distinct degrees admits the expansion, $D_N = BN^{1/3} + C + \dots$ for strictly linear preferential attachment. This allows us to extract a precise estimate of B from the data that is in excellent agreement with equation (5).

The general behavior outlined above for the number of distinct degrees and related quantities is also observed in the citation network of the Physical Review. Because this




Figure 4. The maximum degree (number of citations) and the number of distinct degrees for the Physical Review citation network during the period 1893–2003. Also shown are the locations of the last dimer and the first hole. The dashed lines are the power-law fits with respective exponents 0.849, 0.627, 0.474, and 0.430. The data are measured at 20 equally-spaced network sizes as discussed in the text.

journal has grown roughly exponentially with time [20, 21], it is not appropriate to use publication date as a proxy for the network size. Since the citation data is presented as a list of links, each in the form of *citing paper* \rightarrow *cited paper*, it is more natural to use the chronologically-ordered number of links as the proxy for network size. We use the Physical Review citation data as of 2003, which contain $L = 3,110,866$ total links (citations). The maximum network degree (the highest-cited paper), the location of the last dimer, the number of distinct degrees, and the location of the first hole dimer are measured when the network size is $Lm/20$, with $m = 1, 2, \dots, 20$ (figure 4).

Naive power-law fits to the first three datasets in figure 4 give $k_{\max} \sim L^{0.849}$, $D_L \sim L^{0.627}$, and $\langle h_1 \rangle \sim L^{0.474}$. Let us provisionally assume that the citation distribution has a power-law dependence on L and, by implication, the same dependence on N .³ Using $k_{\max} \sim N^{1/(\nu-1)}$ and the dependences for the number of distinct degrees and location of the last dimer given in equations (3) and (18), we infer the respective exponents for the degree distribution exponent values of 2.18, 2.09, and 2.11. Thus these three properties are internally consistent under the assumption the citation distribution has a power-law form with exponent in the range 2.1–2.2.

Our simulation results indicate that the random quantity D_N is self-averaging. For strictly linear preferential attachment, we find that the standard deviation grows as $\sqrt{\langle D_N^2 \rangle - \langle D_N \rangle^2} \sim N^{1/6}$. Moreover, the probability distribution $\Pi(D_N)$ of distinct degrees fits the Gaussian

$$\Pi(D_N) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(D_N - \langle D_N \rangle)^2 / 2\sigma^2} \quad (6)$$

³ While a power-law gives a reasonable visual fit to the data, later and larger-scale analyses [20], [22]–[25] suggest that the citation distribution has a log-normal or stretched exponential behavior, rather than a power-law form.




Figure 5. Scaled distribution of distinct degrees $f(z)$ for N up to 10^7 , with 10^4 network realizations for each N , for redirection probability $\frac{1}{2}$ (corresponding to strictly linear preferential attachment). The smooth curve is a Gaussian fit to the data. Visually identical data occurs for other redirection probabilities.

extremely well (figure 5). In appropriately scaled coordinates, this form universally holds for any redirection probability (equivalently different λ values in the attachment rate $A_k = k + \lambda$). Moreover, the scaled distributions $f(z) \equiv \sqrt{2\pi\sigma^2} \Pi(D_N)$, with $z = \sqrt{(\langle D_N^2 \rangle - \langle D_N \rangle^2)/2\sigma^2}$, are virtually identical for different λ values.

3. The first hole

We now study properties of the degree distribution in the sparse regime, where not every degree is represented. First consider the location of the first ‘hole’ in the degree distribution—the smallest degree value for which $N_k = 0$. We define h_1 as the degree value of the first hole, h_2 as the degree of the second hole, etc.

To determine the location of the first hole, it is useful to use the probability $P(h)$ that there are no holes in the degree distribution within the range $[1, h]$. This coincides with the probability that there is at least one node of degree k for every k between 1 and h . Again under the assumption that the number of nodes of degree k is given by an independent Poisson distribution for each k , this probability is given by

$$P(h) = \prod_{1 \leq k \leq h} [1 - e^{-N_k}]. \quad (7)$$

We estimate the location of the first hole from the criterion $P(h_1) = \frac{1}{2}$; however, any constant between 0 and 1 could equally well be chosen in this condition. Taking the logarithm of (7) and using $\ln [1 - e^{-N_k}] \approx -e^{-N_k}$ (which is justifiable since $e^{-N_k} \ll 1$ when $k \leq h_1$), gives the following for the average location $\langle h_1 \rangle$ of the first hole:

$$\int_1^{\langle h_1 \rangle} dk e^{-N_k} = \ln 2. \quad (8)$$

Using equation (2) in (8) we find

$$\langle h_1 \rangle \simeq \left(\frac{\nu NR}{\ln H} \right)^{1/\nu}, \quad H \sim \frac{\nu NR}{[\ln(\nu NR)]^{1+\nu}}. \quad (9a)$$

Since H appears inside the logarithm, one can ignore the logarithmic factor in H itself, thereby giving the simpler and still asymptotically exact formula

$$\langle h_1 \rangle \simeq \left(\frac{\nu NR}{\ln N} \right)^{1/\nu}. \quad (9b)$$

It is worth noting that the naive calculation that leads to (3) for the number of distinct degrees ignores the possibility that holes exist in the range $k \leq (NR)^{1/\nu}$. According to equation (9b), however, the first hole appears earlier than $(NR)^{1/\nu}$ in the $N \rightarrow \infty$ limit. For a terrestrial-scale network with, say $N = 10^9$ nodes, the location of the first hole will be roughly three times smaller than that predicted by the naive estimate (3).

4. Last doublet and last dimer

Somewhere in the tail of the degree distribution lie the last doublet, the largest two consecutive k values for which $N_k > 0$, and the last dimer, the largest k value for which $N_k = 2$ (figure 2). Starting with degree 1, the degree distribution first consists of a long string of consecutive ‘occupied’ degrees $1 \leq k < h_1$, followed by a second string in the degree range $h_1 < k < h_2$, etc. As the degree increases, these strings become progressively shorter and above a certain threshold all remaining strings are singlets. For a large network, the last string that is not a singlet will almost certainly be a doublet (with probability approaching 1 as $N \rightarrow \infty$). We now determine the average position of this last doublet.

The probability to have a doublet at $(k, k+1)$ is N_k^2 when $k \gg K = (NR)^{1/\nu}$. To estimate the position of the last doublet $(\delta, \delta+1)$ we employ the extremal criterion

$$\sum_{k \geq \delta} N_k^2 \sim 1 \quad (10)$$

that there should be of the order of one doublet in the degree range (δ, ∞) . Using $N_k \sim NRk^{-\nu}$, we obtain

$$\langle \delta \rangle = C(RN)^{1/(\nu-1/2)}, \quad (11a)$$

with C a constant, for the average position of the last doublet. Notice that the position of the last doublet also coincides, up to a prefactor of the order of 1, to the position of the last dimer. A more precise approach to determine the average location of the last doublet gives the amplitude as

$$C = (2\nu - 1)^{-1/(2\nu-1)} \Gamma\left(\frac{2\nu - 2}{2\nu - 1}\right). \quad (11b)$$

To establish (11b), we use the independent Poisson approximation to write, for the probability $F(\delta)$ to have no doublets in the degree range $k > \delta$,

$$F(\delta) = \prod_{k > \delta} \left[1 - (1 - e^{-N_k})^2 \right]. \quad (12)$$

This expression is the straightforward generalization of equation (7) to the case of dimers. Since the average number of nodes with degrees in the range $k > \delta$ is small, the product on the right-hand side of (12) simplifies to $\exp[-\int_{\delta}^{\infty} dk N_k^2]$. Computing the integral gives

$$F = \exp[-(\delta_0/\delta)^{2\nu-1}], \quad \delta_0 = \left(\frac{R^2 N^2}{2\nu - 1} \right)^{1/(2\nu-1)}. \quad (13)$$

The probability density $\Phi = dF/d\delta$ for the last doublet is then

$$\Phi(\delta) = \frac{2\nu - 1}{\delta} \left(\frac{\delta_0}{\delta} \right)^{2\nu-1} \exp[-(\delta_0/\delta)^{2\nu-1}], \quad (14)$$

from which the average position of the last doublet is given by

$$\langle \delta \rangle = \int_0^{\infty} d\delta \delta \Phi(\delta) = \int_0^{\infty} d\delta [1 - F(\delta)]. \quad (15a)$$

Substituting (13) into (15a) leads to

$$\langle \delta \rangle = \Gamma \left(\frac{2\nu - 2}{2\nu - 1} \right) \delta_0, \quad (15b)$$

which reproduces (11). Similarly, the mean-square position of the last doublet is

$$\langle \delta^2 \rangle = \int_0^{\infty} d\delta \delta^2 \Phi(\delta) = 2 \int_0^{\infty} d\delta \delta [1 - F(\delta)], \quad (16)$$

from which the variance is

$$\langle \delta^2 \rangle - \langle \delta \rangle^2 = \left[\Gamma \left(\frac{2\nu - 3}{2\nu - 1} \right) - \Gamma^2 \left(\frac{2\nu - 2}{2\nu - 1} \right) \right] \delta_0^2. \quad (17)$$

For strictly linear preferential attachment network growth, the above results reduce to

$$\begin{aligned} \langle \delta \rangle &= AN^{2/5}, & A &= \left(\frac{16}{5} \right)^{1/5} \Gamma \left(\frac{4}{5} \right) = 1.469 158 \dots \\ \langle \delta^2 \rangle^{1/2} &= V \langle \delta \rangle, & V &= \frac{\sqrt{\Gamma(3/5)}}{\Gamma(4/5)} = 1.048 182 \dots \end{aligned} \quad (18)$$

Following the same line of reasoning, the position of the last triplet, $(\tau - 1, \tau, \tau + 1)$, is given by

$$\langle \tau \rangle \sim N^{1/(\nu-1/3)}. \quad (19)$$

For strictly linear preferential attachment, this result gives the dependence $\langle \tau \rangle \sim N^{3/8}$. Our simulation data are consistent with the predictions (18)–(19) (figure 6).

5. Discussion

For any broadly distributed integer-valued variable, the underlying distribution exhibits intriguing features that stem from the combined influences of discreteness and finiteness. Such a distribution is smooth in a dense regime, where every integer value of the variable has a non-zero probability of occurrence. In the complementary sparse regime, a variety of statistical anomalies arise that quantify the extent of the sparseness (figure 2).




Figure 6. The location of the last hole and the last doublet as a function of N for 10^4 realizations of networks that are grown by strictly linear preferential attachment. The dashed curve is the prediction (9a), while the straight dashed line is the prediction from equation (18).

For the degree distribution of complex networks that generically have power-law tails, $N_k \sim N/k^\nu$, our main results are as follows. (i) The number of distinct degrees in a network of N nodes scales as $N^{1/\nu}$. This generic behavior is also observed in the citation network of the Physical Review. (ii) The distribution in the number of distinct degrees is very well fitted by a universal Gaussian function. (iii) There is a rich set of behaviors for basic characteristics of the sparse regime, such as the positions of holes (zeros) in the distribution, as well as the locations of doublets, triplets, etc., and the locations of dimers, trimers, etc. All of these quantities can be determined by simple probabilistic reasoning.

Our analysis tacitly assumed that the number of nodes of different degrees, N_i and N_j for $i \neq j$, are uncorrelated, and that the N_k s are Poisson distributed random quantities. While these assumptions are questionable in the sparse regime, predictions that are based on these assumptions are in excellent agreement with results from simulations of preferential attachment networks. While we believe that our predictions are asymptotically exact, a more rigorous analysis is needed to justify them and explain their validity (or at least their impressive accuracy). A challenging extension of this work is to probe the fluctuations in the total number of distinct degrees. The mechanism for the observed Gaussian shape of the distribution of distinct degrees is not at all evident. In fact, for networks that grow by redirection, with redirection becoming more certain as the degree of the ancestor node increases, the total number of distinct degrees is not even a self-averaging quantity [26].

Our methods apply equally well to other heavy-tailed integer-valued distributions, such as the cluster-size distribution in classical percolation [27] and in protein interaction and regulatory networks [28]. The latter models often exhibit an infinite-order percolation transition, in which the cluster-size distribution has an algebraic tail in the entire non-percolating phase [29]–[36]. Our approach leads to new results for the total number of

distinct cluster types C_N , for the position of the first hole (the minimal size that is not present), etc.

For concreteness, consider networks that are built by adding nodes one at a time with each new node connecting to k randomly chosen existing nodes with probability p_k [35, 36]. While the set of probabilities $p_k, k = 0, 1, 2, \dots$, with $\sum_{k \geq 0} p_k = 1$, fully defines the network ensemble, only the first two moments, $\langle k \rangle = \sum_{k \geq 1} k p_k$ and $\Delta = \langle k^2 \rangle - \langle k \rangle^2$, matter in determining large-scale properties. In the non-percolating phase, $\langle k \rangle < \frac{1}{2}$ and $\Delta < \frac{1}{4}$, we use the decay exponent for the cluster-size distribution that was determined in [36] to obtain

$$C_N \sim N^\delta, \quad \delta = \frac{1 - \sqrt{1 - 4\Delta}}{3 - \sqrt{1 - 4\Delta}}.$$

At the percolation transition, $\langle k \rangle < \frac{1}{2}$ and $\Delta = \frac{1}{4}$, the tail of the cluster-size distribution contains universal (independent of $\langle k \rangle$ and Δ) algebraic and logarithmic factors, namely $c_s \simeq 2(1 - 2\langle k \rangle)^{-2} s^{-3} (\ln s)^{-2}$ for $s \gg 1$. A straightforward generalization of our previous analysis shows that the total number of distinct cluster types grows as

$$C_N \simeq 2^{1/3} \Gamma\left(\frac{2}{3}\right) \left(\frac{3}{1 - 2\langle k \rangle}\right)^{2/3} \left[\frac{N}{(\ln N)^2}\right]^{1/3}.$$

As a final note, this work has focused broadly on properties associated with the support of discrete distribution. The averages of these properties over a large ensemble of networks have systematic dependences on the number of nodes N in the network; however, the behavior in each network realization may not be monotonic. Thus while k_{\max} is clearly a non-decreasing function of N , the number of distinct degrees and the locations of quantities like the first hole or the last doublet can both increase or decrease with N . This intriguing aspect of the problem may provide a more detailed understanding of how a complex network actually grows.

Acknowledgments

We gratefully acknowledge partial financial support from AFOSR and DARPA grant #FA9550-12-1-0391, and from NSF grant No. DMR-1205797. We also thank A Gabel for collaborations at a preliminary stage of this project. Finally, we thank the American Physical Society editorial office for providing the citation data.

References

- [1] Newman M E J, 2010 *Networks: An Introduction* (Oxford: Oxford University Press)
- [2] Albert R and Barabási A-L, 2002 *Rev. Mod. Phys.* **74** 47
- [3] Yule G U, 1925 *Phil. Trans. R. Soc. B* **213** 21
- [4] Simon H A, 1955 *Biometrika* **42** 425
- [5] de Solla Price D J, 1965 *Science* **149** 510
- [6] de Solla Price D J, 1976 *J. Am. Soc. Inform. Sci.* **27** 292
- [7] Barabási A L and Albert R, 1999 *Science* **286** 509
- [8] Krapivsky P L, Redner S and Leyvraz F, 2000 *Phys. Rev. Lett.* **85** 4629
- [9] Dorogovtsev S N, Mendes J F F and Samukhin A N, 2000 *Phys. Rev. Lett.* **85** 4633
- [10] Herdan G, 1960 *Type-Token Mathematics* (The Hague: Mouton)
- [11] Heaps H S, 1978 *Information Retrieval: Computational and Theoretical Aspects* (New York: Academic)

- [11] van Leijenhorst D D and van der Weide T P, 2005 *Inf. Sci.* **170** 263
- [12] Lu L, Zhang Z-K and Zhou T, 2010 *PLoS One* **5** e14139
- [13] Gerlach M and Altmann E G, 2012 arXiv:1212.1362
- [14] Zipf G K, 1936 *The Psychobiology of Language* (London: Routledge)
- [15] Erdős P and Gallai T, 1960 *Matematikai Lapok* **11** 264
- [16] Del Genio C I, Kim H, Toroczkai Z and Bassler K E, 2010 *PLoS One* **5** e10012
- [17] Del Genio C I, Gross T and Bassler K E, 2011 *Phys. Rev. Lett.* **107** 178701
- [18] Krapivsky P L and Redner S, 2001 *Phys. Rev. E* **63** 066123
- [19] Krapivsky P L and Redner S, 2002 *J. Phys. A: Math. Theor.* **35** 9517
- [20] Redner S, 2005 *Phys. Today* **58** 49
Redner S, 2004 arXiv:physics/0407137
- [21] Krapivsky P L and Redner S, 2005 *Phys. Rev. E* **71** 036118
- [22] Wallace M L, Lariviere V and Gingras Y, 2009 *J. Informetrics* **3** 296
- [23] Phillips J C, 2009 arxiv:0909.5363
- [24] Peterson G J, Presse S and Dill K A, 2010 *Proc. Nat. Acad. Sci.* **107** 16023
- [25] Golosovsky M and Solomon S, 2012 *Phys. Rev. Lett.* **109** 098701
Golosovsky M and Solomon S, 2012 *Eur. Phys. J. Spec. Top.* **205** 303
- [26] Gabel A, Krapivsky P L and Redner S, in preparation
- [27] Stauffer D and Aharony A, 1991 *Introduction to Percolation Theory* 2nd edn (London: Taylor and Francis)
- [28] Uetz P L *et al*, 2000 *Nature* **403** 623
Ito T *et al*, 2000 *Proc. Nat. Acad. Sci.* **97** 1143
- [29] Callaway D S, Hopcroft J E, Kleinberg J M, Newman M E J and Strogatz S H, 2001 *Phys. Rev. E* **64** 041902
- [30] Dorogovtsev S N, Mendes J F F and Samukhin A N, 2001 *Phys. Rev. E* **64** 066110
- [31] Krapivsky P L and Redner S, 2002 *Comput. Netw.* **39** 261
- [32] Kim J, Krapivsky P L, Kahng B and Redner S, 2002 *Phys. Rev. E* **66** 055102
- [33] Lancaster D, 2002 *J. Phys. A: Math. Theor.* **35** 1179
- [34] Bauer M and Bernard D, 2003 *J. Stat. Phys.* **111** 703
- [35] Coulomb C and Bauer M, 2003 *Eur. Phys. J. B* **35** 377
- [36] Krapivsky P L and Derrida B, 2004 *Physica A* **340** 714