

# Supersymmetry (SUSY) Past, Present and Future

Krzysztof Sliwa



Tufts University  
Department of Physics and Astronomy  
Medford, Massachusetts 02155, USA

NEPPSR Craigville, August 19, 2005

# symmetries $\Leftrightarrow$ conservation laws

- Emmy Noether discovered the connection between symmetries and conservation laws while working with David Hilbert and Felix Klein in Gottingen
- In 1918 she proved two theorems, for finite continuous groups and infinite continuous groups which are the foundations of the modern (XXth century) physics. The theorems are collectively known as “Noether’s theorem”
- Informally, Noether’s theorem says:

differentiable symmetry generated by local actions  $\Leftrightarrow$  conserved current

or

there is one-to-one correspondence between symmetries and conservation laws  
symmetry  $\Leftrightarrow$  conservation law

# symmetries $\Leftrightarrow$ conservation laws

- examples:
  - **energy** is conserved if and only if (iff) the physical laws are invariant under **time translations** (if the form of physics laws do not depend on time)
  - **linear momentum** is conserved only iff the physical laws are invariant under **space translations** (if the form of physics laws do not depend on the position)
  - **angular momentum** is conserved iff the physical laws are invariant under **rotations** (if the physics laws do not depend on orientation; if only true about a particular direction  $\Leftrightarrow$  only the component of angular momentum in that direction is conserved)

# symmetries $\Leftrightarrow$ conservation laws

- Symmetries observed in physics:
  - Symmetries of discrete space-time transformations: parity, time-reversal, charge conjugation
  - Symmetries of continuous space-time transformations: translational and rotational invariance and Lorentz (space-time rotations) invariance
  - Symmetries of permutations: lead to two kind of particles: bosons, which obey Bose-Einstein statistics, and fermions, which obey Fermi-Dirac statistics
  - Gauge symmetries: internal symmetries inherent from the nature of the field associated with a given particle carrying such attributes as electric charge - U(1), color - SU(3) et cetera (**conservation of electric charge**  $\Leftrightarrow$  invariance under the global **phase transformation** in the **internal space**; electromagnetic field  $\Leftrightarrow$  invariance under the local phase transformation; et cetera....you'll learn all this in the next 2 years!)

# symmetries $\Leftrightarrow$ conservation laws

- Modern particle physics is based entirely on the idea of underlying internal symmetries:
  - The electro-weak sector is based upon the (internal) symmetries which the electromagnetic and weak interactions obey -  $U(1)$  and  $SU(2)$
  - The strong sector of the Standard Model (SM), quantum chromodynamics (QCD) is based on the (internal)  $SU(3)$  symmetries observed in hadron spectroscopy
  - Spontaneous symmetry breaking has been proposed to explain massive weak bosons ( $Z$ ,  $W$ ) and the massless photon. The prediction of the  $W$  and  $Z$  bosons came from symmetry arguments and the discovery of these particles at CERN was one of the greatest successes of modern particle physics

# STANDARD MODEL


- Current understanding of elementary particles and their strong and electro-weak interactions is given by Standard Model, a gauge theory based on the following internal symmetries:

$$SU(3)_c \times SU(2)_I \times U(1)_Y$$

- The  $SU(3)$  is an unbroken symmetry, it gives QCD, a quantum theory of strong interactions, whose carriers (gluons) are massless
- $SU(2) \times U(1)$  (quantum theory of electroweak interactions) is spontaneously broken by the Higgs mechanism; which gives mass to electroweak bosons ( $W^+$ ,  $W^-$ ,  $Z^0$  and a massless photon)
- In the Minimal Standard Model, the Higgs sector is the simplest possible: contains two complex Higgs fields, which after giving masses to  $W, Z$  give leaves a **neutral scalar Higgs particle which should be observed** - the ONLY particle not yet discovered in MSM

# STANDARD MODEL

- Matter is build of fermions - quarks and leptons, three families of each, with corresponding antiparticles; quarks come in three colors
- Bosons are carriers of interactions: 8 massless gluons, 3 heavy weak bosons (W,Z) and 1 massless photon
- A massive scalar Higgs field permeates the Universe and is (in some way) responsible for masses of other particles



~28 parameters NOT predicted by SM:



- masses of 6 quarks
- masses of 6 leptons
- coupling constants of SU(3), SU(2) and U(1)
- Higgs mass and vacuum expectation value
- Cabibbo-Kobayashi-Maskawa matrix angles and complex phase
- Maki-Nakagawa-Sakata matrix angles and complex phase
- QCD phase  $\theta$

**ALL MUST BE MEASURED !!!**

# STANDARD MODEL

- **Masses** of quarks and leptons, as well as those of carriers of interactions and Higgs scalar particle are fundamental parameters of SM - **to be determined by measurement**
- **mixing angles** in quark and lepton sector, and the **phases** are also parameters **to be measured**
- It is possible to verify the **internal consistency of SM** through precise measurements: together with other already very precise EW measurements, precise measurements of W and top mass constrain Higgs mass. Fundamental consistency tests of Standard Model; sensitivity through radiative corrections (quadratic in  $m_t$ , logarithmic in  $m_H$ )

**COMPARE WITH DIRECT LIMITS ON HIGGS MASS**






# spontaneous breaking of the electroweak symmetry by Higgs mechanism

- This part of SM is the only remaining untested part of SM. Higgs has not been observed as of yet; remember, the EW symmetry could be broken in a different way, not necessarily like in MSM
- Difficulties with the elementary Higgs sector: suppose that SM is just an effective theory and that NEW physics is at some scale  $\Lambda$ .

the quantum corrections to fermion masses would depend only logarithmically on scale  $\Lambda$ :




$$\delta m_f \sim m_f \ln \Lambda$$

# spontaneous breaking of the electroweak symmetry by Higgs mechanism

- Difficulties with the elementary Higgs sector: the analogous quantum corrections to scalar particle (Higgs) would exhibit a quadratic dependence on scale  $\Lambda$ . This means that Higgs mass is VERY sensitive to the scale of the NEW physics => FINE TUNING PROBLEM (for  $m_0$ ) as  $m_H = O(100)$  GeV in SM !!

$$m_H^2 = -m_0^2 + g^2\Lambda^2$$

SM cannot be valid for very large momenta, the scale  $\Lambda$  serves as a cutoff above which physics not contained in SM becomes important. At least one such scale, Planck scale at which gravity becomes relevant,  $\Lambda = O(10^{19})$  GeV, must be present in any theory.



$$\delta m_H^2 \sim \Lambda^2$$


# spontaneous breaking of the electroweak symmetry by Higgs mechanism

- This fine tuning has to be performed for each order of perturbation theory; this is a very unpleasant feature of SM
- This sensitivity is called also the GAUGE HIERARCHY PROBLEM, as the Higgs mass is related to the weak boson masses in the spontaneously broken gauge theory. One may say that the original problem of how to give masses to weak gauge bosons in a gauge invariant way was only partially solved by Higgs mechanism, and the problem was transferred to a new level, where the new puzzle is how to keep Higgs mass stable against large quantum corrections from the higher energy scales
- A method of controlling Higgs mass divergence other than fine tuning of parameters would be very welcomed

# supersymmetry - the most elegant solution?

- the interesting thing about the scalar mass divergencies from virtual particle loops (quantum corrections) is that *virtual fermions and virtual bosons contribute with opposite signs and would cancel each other exactly if for every boson there was a fermion of the same mass and charge - divergencies would cancel without any fine tuning and in all orders of perturbation theory !!*
- supersymmetry is such a symmetry: it connects bosons to fermions, it introduces a fermionic partner to every boson and vice-versa, identical in all quantum numbers; such boson  $\Leftrightarrow$  fermion connection is unique to supersymmetry; all the symmetries listed before provide no such connection

# Supersymmetry



# supersymmetry - the most elegant solution?

- at the quantum mechanical level, this Fermi-Bose symmetry would require some quantum operator,  $Q$ , whose action would be to transform bosons into fermions and vice-versa

$$Q|\text{fermion}\rangle = |\text{boson}\rangle$$

$$Q|\text{boson}\rangle = |\text{fermion}\rangle$$

- and since this is a symmetry, this operator must commute with the Hamiltonian

$$[Q, H] = 0$$

# supersymmetry - the most elegant solution?

- Such a theory is called a *supersymmetric theory* and the *operator Q* is called the *supercharge*. Since the operator Q changes a particle with spin  $1/2$  to a particle with spin  $1$  or  $0$ , the Q itself must be a spinor that carries spin  $1/2$  of its own
- Bosons are particles with integer spins, they obey Bose-Einstein statistics, any number of them may occupy the same quantum state at a time. Fermions carry half-integer spin (or odd multiples of  $1/2$ ), they obey Fermi statistics and only one fermion can occupy any given quantum state at a time. The classical limit of quantum mechanics is approached when the occupation numbers of available states are very high. The quantum photon field behaves like the classical EM field described by Maxwell's equations. However, there is NO classical limit for fermions, fermion fields are quantum phenomena.
- A symmetry that interchanges fermions and bosons is a symmetry that exchanges physics that has a classical limit with physics with NO classical limit - POTENTIALLY EXTREMELY POWERFUL and interesting

# supersymmetry - the most elegant solution?

- Obviously, if supersymmetry were real, it must be somehow broken as we have not yet observed superparticles. One needs to allow such breaking of supersymmetry while still keeping the ability of such a theory to solve the gauge hierarchy problem. Not easy, depends on the scale at which SUSY is broken, and on how it is broken. To some extent it remains still an open question
- Another reason for SUSY theories being attractive is that in string theories the most viable versions are supersymmetric
- Local supersymmetry could also be a viable theory of gravity, supergravity.



# supersymmetry - the most elegant solution?

- space-time symmetries in relativistic QM are contained in the Poincare group, it includes symmetries under spatial rotations, translations in space and time and space-time boosts (space-time rotations)
- a symmetry group is described by the algebra of the group which is defined by a set of commutation relations. For the Poincare group:

$$[P^\mu, P^\nu] = 0$$

$$[J^{\mu\nu}, P^\kappa] = P^\mu \eta^{\nu\kappa} - P^\nu \eta^{\mu\kappa}$$

$$[J^{\mu\nu}, J^{\kappa\lambda}] = J^{\mu\lambda} \eta^{\nu\kappa} - J^{\nu\lambda} \eta^{\mu\kappa} - J^{\mu\kappa} \eta^{\nu\lambda} + J^{\nu\kappa} \eta^{\mu\lambda}$$

$P^\mu$  is the momentum generator which generates space and time translations, the Lorentz matrices  $J^{\nu\kappa}$  generate rotations in space and Lorentz boosts (rotations) in space-time, and  $\eta^{\mu\kappa}$  is the metric tensor.

These are all bosonic symmetries, which should be true as energy, momentum and angular momentum conservation and Lorentz invariance are present in classical physics

# supersymmetry - the most elegant solution?

- our world is described by:

Poincare (space-time) symmetry: with generators  $P^\mu, J^{\nu\kappa}$

internal symmetries ( $U(1)\times SU(2)\times SU(3)$  of SM): with generators  $T_a$

- However, the Poincare group also has representations that describe fermions. This should be expected as spin 1/2 particles appear as solutions to a relativistically invariant equation - the Dirac equation. If there exist spin 1/2 particles could there be spin 1/2 symmetry generators in a space-time symmetry algebra?
- This would be an extension of Poincare group of symmetries valid for relativistic QFT in  $D=4$

# supersymmetry - the most elegant solution?

- in 1971 Golfand and Likhtman (whose work was forgotten for years..):

$$[P^\mu, P^\nu] = 0$$

$$[P^\mu, Q_a] = [P^\mu, \bar{Q}_a] = 0$$

$$\{Q_a, Q_b\} = \{\bar{Q}_a, \bar{Q}_b\} = 0$$

$$\{Q_a, \bar{Q}_b\} = 2\gamma_{ab}^\mu P_\mu$$

Note  $E = H = P^0 \Rightarrow [Q_a, H] = 0 \Rightarrow Q$  is a conserved charge.

$Q_a$  is fermionic generator (spinor) with  $\bar{Q}_b$  its complex conjugate. What are these new symmetry generators  $Q$ ? These are the *supercharges* mentioned before (note anticommutators  $\{, \}$  instead of commutators  $[, ]$ )

- If there is just one fermionic generator (supercharge)  $Q$  we call such a theory N=1 SUSY; if there are two, we have N=2 SUSY, et cetera...
- in 1974 Wess and Zumino wrote a Lagrangian with the same symmetries

# SM and MSSM particle spectrum

Standard Model Particles		SUSY Partners		
Particles	States	Sparticles	States	Mixtures
quarks ( $q$ ) (spin- $\frac{1}{2}$ )	$\begin{pmatrix} u \\ d \end{pmatrix}_L, u_R, d_R$ $\begin{pmatrix} c \\ s \end{pmatrix}_L, c_R, s_R$ $\begin{pmatrix} t \\ b \end{pmatrix}_L, t_R, b_R$	squarks ( $\tilde{q}$ ) (spin-0)	$\begin{pmatrix} \tilde{u} \\ \tilde{d} \end{pmatrix}_L, \tilde{u}_R, \tilde{d}_R$ $\begin{pmatrix} \tilde{c} \\ \tilde{s} \end{pmatrix}_L, \tilde{c}_R, \tilde{s}_R$ $\begin{pmatrix} \tilde{t} \\ \tilde{b} \end{pmatrix}_L, \tilde{t}_R, \tilde{b}_R$	$\tilde{t}_{1,2}, \tilde{b}_{1,2}$
leptons ( $l$ ) (spin- $\frac{1}{2}$ )	$\begin{pmatrix} e \\ \nu_e \end{pmatrix}_L, e_R$ $\begin{pmatrix} \mu \\ \nu_\mu \end{pmatrix}_L, \mu_R$ $\begin{pmatrix} \tau \\ \nu_\tau \end{pmatrix}_L, \tau_R$	sleptons ( $\tilde{l}$ ) (spin-0)	$\begin{pmatrix} \tilde{e} \\ \tilde{\nu}_e \end{pmatrix}_L, \tilde{e}_R$ $\begin{pmatrix} \tilde{\mu} \\ \tilde{\nu}_\mu \end{pmatrix}_L, \tilde{\mu}_R$ $\begin{pmatrix} \tilde{\tau} \\ \tilde{\nu}_\tau \end{pmatrix}_L, \tilde{\tau}_R$	$\tilde{\tau}_{1,2}$
gauge/Higgs bosons (spin-1, spin-0)	$g, Z, \gamma, h, H, A$ $W^\pm, H^\pm$	gauginos/Higgsinos (spin- $\frac{1}{2}$ )	$\tilde{g}, \tilde{Z}, \tilde{\gamma}, \tilde{H}_1^0$ } $\tilde{W}^\pm, \tilde{H}^\pm$ }	$\tilde{\chi}_{1,2,3,4}^0$ $\tilde{\chi}_{1,2}^\pm$
graviton (spin-2)	$G$	gravitino (spin- $\frac{3}{2}$ )	$\tilde{G}$	

# supersymmetry - classification of models

- in the past 30 years extended studies of low-energy SUSY and a number of experimental searches => no evidence for SUSY. However, if SUSY is broken on a scale of  $\sim 1$  TeV, LHC will have a great chance to discover superparticles
- **the minimal supersymmetric standard model (MSSM)**, two Higgs doublets. Has one less parameter than SM if SUSY is unbroken. Obviously this is not true; supersymmetry must be broken *without* destroying the cancellations which solve the fine-tuning problem => soft SUSY breaking, however no particular way that it is done is assumed. In MSSM R parity is conserved (R=+1 for SM particles, R=-1 for superparticles) which means SUSY particles must be produced in pairs and that the lightest SUSY particle (LSP) is stable (a candidate for “dark matter”)

$$R = (-1)^{3B+2S+L}$$

Difficult to use MSSM for experimental studies because of large number of parameters ( $\sim 105$  free parameters !)


# supersymmetry - classification of models

- Different models of SUSY breaking are used to reduce the number of parameters. They all have a common feature: SUSY is broken in some hidden sector and then transmitted to the MSSM fields. The models differ in how this is done:
- **SUGRA**: in supergravity models all scalar masses ( $M_0$ ), the gaugino mass ( $M_{1/2}$ ), and the A and B parameters are assumed to be unified at at GUT scale ( $\sim 10^{15}$  GeV). Five parameters:  $M_0$ ,  $M_{1/2}$ , A,  $\text{sgn}(\mu)$  and  $\tan\beta$  completely determine the mass spectrum and decay patterns of particles ( $\tan\beta$ =ratio of vacuum expectation values of the two Higgs doublets, A-trilinear coupling and  $\text{sgn}(\mu)$ -sign of supersymmetric Higgs parameter. Mediating interaction is gravitational

# supersymmetry - classification of models


- **GMSB**: (gauge mediated symmetry breaking) rather than using gravity to transmit the SUSY breaking, gauge interactions are used. The messenger sector consists of some particles,  $X$ , which have SM interactions and are aware of SUSY breaking. The LSP is almost massless gravitino. The model has 6 parameters....
- **AMSB**: (anomaly mediated symmetry breaking); the mAMSB model has 5 parameters, very similar to mSUGRA
- as you can imagine, many others.....

# spectrum of particle masses in SUSY models







# Merging forces



# running coupling constants in SM and MSSM models



# RUN-II AT TEVATRON



Fermi National Accelerator Laboratory

# RUN-II AT TEVATRON

## 2001-?

New Main Injector  $\Rightarrow$  CM energy ( $\sqrt{s}$ ) increased from 1800 GeV to 1960 GeV ( $t\bar{t}$  cross section increases by  $\sim 35\%$ )

Different beam crossing time (396 ns and 132 ns later (?), instead of  $3.5 \mu\text{s}$  in Run-I) - fewer multiple interactions


Significant upgrades to both detectors:

D0 : addition of SVX to allow better b-tagging  
addition of a solenoid to allow track momentum reconstruction


CDF : new calorimeter for  $1.1 < |\eta| < 3.5$  (much better energy resolution)  
new (longer) SVX with double the Run-I tagging efficiency

# RUN-II AT TEVATRON 2001-?

- CDF and D0: well-understood, mature detectors with excellent particle identification, coverage, tracking and triggering





# RUN-II AT TEVATRON 2001 - ?



D0 detector in its current configuration

# RUN-II AT TEVATRON 2001 - ?



CDF detector in its current configuration

## chargino-neutralino searches

$$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow l^\pm l^\mp l^\pm \tilde{\chi}_1^0 \tilde{\chi}_1^0 X$$

- at Tevatron: look for lightest chargino, 2<sup>nd</sup> neutralino
- final state with many leptons, large  $E_T^{Miss}$  from LSP
- one of the SUSY “golden modes”
  - small SM backgrounds but small (EW) cross sections
  - striking signature





# chargino-neutralino searches (D0)

In models with  $R_p$ :

3 leptons +  $E_T^{Miss}$

- ➔  $\sigma_{BR} \sim 0.2$  pb
- ➔ Very clean signature
- ➔ SM background very small

Selection	Bkgn'd expected	data
ee+l	$0.21 \pm 0.12$	0
eμ+l	$0.31 \pm 0.13$	0
μμ+l	$1.75 \pm 0.57$	2
μ <sup>±</sup> μ <sup>±</sup>	$0.64 \pm 0.38$	1
eτ+l	$0.58 \pm 0.14$	0
μτ+l	$0.36 \pm 0.13$	1
<b>SUM</b>	<b><math>3.85 \pm 0.75</math></b>	<b>4</b>



# chargino-neutralino searches (D0)

## “3l-max”

- 

Limits :

➔  $M(\chi_{\pm 1}^{\pm}) > 116 \text{ GeV}/c^2$


## “Heavy Squarks”

- 

➔  $M(\chi_{\pm 1}^{\pm}) > 128 \text{ GeV}/c^2$

## “Large $m_0$ ”

- $M(\tilde{l}) \gg M(\chi_2^0, \chi^{\pm})$
- No sensitivity due to smaller leptonic BR's



# chargino-neutralino searches (CDF)

## SELECTION:

2 electrons+  $l(l=e,\mu) \quad |\eta| < 1$

large  $E_T^{Miss} > 15 \text{ GeV}/c^2$

$15 < M_{\parallel} < 76, > 106 \text{ GeV}/c^2$


$|\Delta\phi| < 160$

$N_{jets}(20 \text{ GeV}) < 2$


Process	
mSugra eel	0.5
Bkgnd Expected	$0.16 \pm 0.07$
OBSERVED	0

krzysztof.sliwa@tufts.edu

## Search for $\chi_2^0 \chi_1^\pm \rightarrow ee+I+X$




## Search for $\chi_2^0 \chi_1^\pm \rightarrow ee+I+X$





NEPPSR, Craigville, ...

# chargino-neutralino searches (D0)



#jets( $P_T$ (GeV))	$\sum P_t^{\text{jet}}$	$E_T^{\text{Miss}}$	Bkgd Expected	data
2 jets(60,50)	250 GeV	175 GeV	$12.8 \pm 5.4$	12
3 jets(60,40,25)	325 GeV	100 GeV	$6.1 \pm 3.1$	5
4 jets(60,40,30,25)	175 GeV	75 GeV	$9.3 \pm 0.5$	10



Limits ( $\tan\beta=3, A_0=0, \mu<0, \tilde{q}=u,d,c,s,b$ ):

→ 2j :  $M_0=25 \text{ GeV} \rightarrow \tilde{M}(q) > 318 \text{ GeV}/c^2$


→ 3j :  $M(g)=M(q) \rightarrow \tilde{M}(q) > 333 \text{ GeV}/c^2$

→ 4j :  $M_0=500 \text{ GeV} \rightarrow M(g) > 233 \text{ GeV}/c^2$

# gluino-sbottom searches (CDF)


$$\tilde{g}\tilde{g} \rightarrow \tilde{b}_1\tilde{b}_1\bar{b}\bar{b} \rightarrow b\bar{b}b\bar{b}\tilde{\chi}_1^0\tilde{\chi}_1^0$$

- striking signature: four b's in final state + large  $E_T^{Miss}$ .
- identify b quark jets to reduce dijet backgrounds
  - use displaced tracks to tag
- efficiency of b-tagging depends on
  - $m(\text{gluino}) - m(\text{sbottom})$
- set limits as function of  $m_{\text{gluino}}, m_{\text{sbottom}}$



$n_{\text{tag}}$	background	observed
=1	$16.4 \pm 3.7$	21
$\geq 2$	$2.6 \pm 0.7$	4

# LHC at CERN: SUSY particles factory?



	$\sqrt{s}$ [TeV]	Luminosity [cm <sup>-2</sup> s <sup>-1</sup> ]	$\int Ldt$ [fb <sup>-1</sup> /y]
Tevatron	2	<10 <sup>32</sup>	0.3
LHC (low lum)	14	10 <sup>33</sup>	10
LHC (high lum)	14	10 <sup>34</sup>	100


Data in 2007?

# Future: SUSY at LHC




0712/mb-26/06/97

## A Toroidal LHC Apparatus




# Future: SUSY at LHC







# SUSY signatures at LHC



- Heavy **gluinos** and **squarks** (strongly interacting particles) produced in initial interaction
- Long decay chains and large mass differences between SUSY states; many **high  $P_T$  objects** are observed (lepton, jets, b-jets)
- **If the model is mSUGRA** R-Parity is conserved, lightest SUSY particle (LSP) is a stable neutralino, cascade decays lead to stable undetected LSP => **large  $E_T^{miss}$  signatures**
- **If the model is GMSB**, LSP is gravitino. Additional signatures from NLSP (next-to-lightest SUSY particle) decays; for example photons (neutralino decays into photon and gravitino) and leptons from slepton decays (from neutralino decaying into lepton and gravitino)
- **If R-parity is not conserved** LSP decays to 3-leptons, 2leptons+1jet, 3 jets;  $E_T^{miss}$  signature is lost

# mSUGRA

- mSUGRA framework: five free parameters:  $m_0$ ,  $m_{1/2}$ ,  $A_0$ ,  $\tan(\beta)$ ,  $\text{sgn}(\mu)$
- sensitivity only weakly dependent on  $A_0$ ,  $\tan(\beta)$ ,  $\text{sgn}(\mu)$
- multiple signatures on most of parameter space:  $E_T^{\text{miss}}$  (dominant signature),  $E_T^{\text{miss}}$  with lepton veto, one lepton, two leptons same sign (SS), two leptons opposite sign (OS)



# mSUGRA: selected points

- **DC1 bulk region point** (new underlying event in generation)
  - $m_0 = 100$  GeV,  $m_{1/2} = 300$  GeV,  $A_0 = -300$  GeV,  $\tan\beta = 6$ ,  $\text{sgn}(\mu) = +$
  - LSP is mostly bino, light  $I_R$  enhance annihilation. 'Bread and butter' region for the LHC experiments
  - $llq$  distributions, tau-tau measurements, third generation squarks (both tau identification and B tagging improved)
- **Coannihilation point**
  - $m_0 = 70$  GeV,  $m_{1/2} = 350$  GeV,  $A_0 = 0$  GeV,  $\tan\beta = 10$ ,  $\text{sgn}(\mu) = +$
  - LSP is pure bino. LSP/sparticle coannihilation. Small slepton-LSP mass difference gives soft leptons in the final state
- **Focus point**
  - $m_0 = 3350$  GeV,  $m_{1/2} = 300$  GeV,  $A_0 = 0$  GeV,  $\tan\beta = 10$ ,  $\text{sgn}(\mu) = +$
  - LSP is Higgsino, near  $\mu^2=0$  bound. Heavy sfermions; all squarks and sleptons have mass  $>2$  TeV, negligible FCNC, CP,  $g_\mu -2$ , etc. Complex events with lots of heavy flavor
- **Funnel region point**
  - $m_0 = 320$  GeV,  $m_{1/2} = 375$  GeV,  $A_0 = 0$  GeV,  $\tan\beta = 50$ ,  $\text{sgn}(\mu) = +$
  - wide H, A for  $\tan\beta \gg 1$  enhance annihilation. Heavy Higgs resonance (funnel); main annihilation chain into  $bb$  pairs
  - dominant tau decays
- **Low mass point** at limit of Tevatron RunII reach
  - $m_0 = 200$  GeV,  $m_{1/2} = 160$  GeV,  $A_0 = -400$  GeV,  $\tan\beta = 10$ ,  $\text{sgn}(\mu) = +$
  - big cross section, but events rather similar to top
  - measure SM processes in presence of SUSY background to show detector is understood

# The $(m_0, m_{1/2})$ - mSUGRA plane

Excluded by  $b \rightarrow s\gamma$   
(CLEO, BELLE)

$m_{1/2} = 114 \text{ GeV}$

$m_{1/2} = 114 \text{ GeV}$

Favored by  $g_\mu - 2$  at the  $2\sigma$  level  
Muon  $g-2$  coll.

WMAP:  $0.094 < \Omega_\chi h^2 < 0.129$

Focus point

$$\tilde{m}_1^0 \approx \tilde{H}$$

Stau1=LSP

Bulk region

t-channel slepton exchange.  
(ATL-PHYS-2004-011)

Stau coannihilation

$$\tilde{m}_1^0 \tilde{q}_1 \rightarrow \tilde{a} \hat{o}$$

Funnel region

$2m_{\tilde{t}_1} \approx m_{H,A}$  s-channel  
Higgs-exchange.

(Ellis et al., Phys. B565 (2003) 176)

# mSUGRA points


The following points in the mSUGRA space have been selected for analysis with the full ATLAS detector simulation (GEANT4).

	$M_0$ (GeV)	$M_{1/2}$ (GeV)	$A_0$	$\tan\beta$	$\text{sgn}(\mu)$	$m_{\text{top}}$ (GeV)
Coannihilation	70	350	0	10	+	175
Focus point	3550	300	0	10	+	175
Funnel region	320	375	0	50	+	175
Bulk (ATL-PHYS-2004-011)	100	300	-300	6	+	175
Scan	130-6000	600,1000	0	10	+	175
low mass point	200	160	-400	10	+	175

Events generated with HERWIG 6.505 (+JIMMY).  
SUSY spectra obtained with ISAJET7.71

**All results shown in this talk are obtained from new full simulation data!**

# DC I: dilepton endpoint





- in mSUGRA R-parity is conserved, all SUSY events contain two neutralinos which escape the detector
- neutralinos are not detected, but one can measure **kinematic endpoints in dilepton invariant mass distributions** rather than mass peaks

$$\tilde{\chi}_2^0 \rightarrow \tilde{l}^{\pm} l^{\mp} \rightarrow \tilde{\chi}_1^0 l^{\pm} l^{\mp}$$

$$M_{ll}^{\max} = \sqrt{\frac{(M_{\tilde{\chi}_2^0}^2 - M_{\tilde{l}}^2)(M_{\tilde{l}}^2 - M_{\tilde{\chi}_1^0}^2)}{M_{\tilde{l}}^2}} = 100.31 \text{ GeV}$$



# coannihilation point:

- Chosen point:  $m_0=70$  GeV;  $m_{1/2}=350$  GeV;  $A_0=0$ ;  $\tan\beta=10$  ;  $\mu>0$ ;
- Small slepton-neutralino mass difference gives soft leptons
- Decays of  $\chi_2^0$  to both  $l_L$  and  $l_R$  kinematically allowed; double dilepton invariant mass edge structure, edges expected at 58 / 98 GeV
- Stau channels enhanced ( $\tan\beta$ ); soft tau signatures, edge expected at 79 GeV. Less clear due to poor tau visible energy resolution



# focus point

- chosen point:  $m_0=3000$  GeV;  $m_{1/2}=215$  GeV;  $A_0=0$ ;  $\tan\beta=10$  ;  $\mu>0$
- large  $m_0 \rightarrow$  sfermions are heavy
- most useful signatures from heavy neutralino decay
- direct three-body decays  $\chi_n^0 \rightarrow \chi_1^0 \ell \ell$
- fit results give:
  - $M(\chi_2^0)-M(\chi_1^0)= 57.45 \pm 0.28$  GeV
  - $M(\chi_3^0)-M(\chi_1^0)= 73.27 \pm 0.47$  GeV





# SM background



Dominant SM background processes:

- ▼ Z+N jets
- ▲ W+N jets
- tt+N jets
- multijets (QCD)
- ▨ sum of all BG

Previous studies are based on Parton shower.

## New SM BG estimation using ME generator (ALPGEN 1.33)

- W/Z + N jets, tt + N jets are generated and processed with the fast ATLAS simulation
- Collinear and soft kinematic regions are assessed with PS (PYTHIA). MLM method used for ME-PS matching.




## SUSY: the “default new physics ??”

- SUSY is perhaps the most explored of “beyond the SM” physics scenarios
- As such, it will perhaps be “blamed” for any deviations from SM physics if observed at Tevatron or at LHC
- The problem will be to prove that, even if a statistically significant deviation from SM predictions is found, the observed events are really due to the supersymmetric particles and NOT to anything else. This will NOT be easy. As you should realize by now, there is an almost continuous spectrum of different SUSY models with different parameters
- Several times in the past (monojets at UA1- see Gary Taubes’s “Nobel Dreams”, CDF- the famous  $e e \gamma \gamma$  event) the excitement ran quite wild about what later proved to be just very rare, but still normal SM, events  
....

# SUSY: the “golden” candidate for “new physics”

- CDF- the famous  $e\bar{e}\gamma\gamma$  MET event: recorded April 28, 1995 in Run-I. Its “a posteriori” probability according to SM  $\sim 10^{-6}$

## $e\bar{e}\gamma\gamma$ $\cancel{E}_T$ Candidate Event



# SUSY: the “golden” candidate for “new physics”

- KEEP YOUR EYES OPEN, LEARN SM WELL, KNOW WHAT TO IS TO BE EXPECTED, EVEN IF IT IS RARE
- DON'T GET TOO EXCITED, MAINTAIN CLARITY OF THOUGHT AT ALL TIMES, IF POSSIBLE
- THE NEXT 5 YEARS COULD BE VERY INTERESTING, TEVATRON AND CERTAINLY LHC WILL PROVIDE A CLOSER LOOK AT THE COMPLETELY UNEXPLORED REGION OF PHASE-SPACE
- REMEMBER THAT THE ATTRACTIVENESS OF SUSY IS REALLY PURELY ESTHETIC, AS IT SOLVES (OR AT LEAST POSTPONES) THE FINE-TUNING PROBLEM AND PROVIDES THE LINK BETWEEN FERMIONS AND BOSONS
- DISCOVERING ANY NEW PHYSICS BEYOND SM WOULD BE A BREAKTROUGH, WE DON'T KNOW WHAT IT WILL BE, IT DOES NOT HAVE TO BE SUSY