

On the Role of Global Warming on the Statistics of Record-Breaking Temperatures

S. Redner^{1,*} and Mark R. Petersen^{2,†}

¹*Center for Polymer Studies & Department of Physics,
Boston University, Boston, Massachusetts, 02215 USA*

²*Computer and Computational Sciences Division and Center for Nonlinear Studies,
Los Alamos National Laboratory, Los Alamos, New Mexico, 87545 USA*

We theoretically study the statistics of record-breaking daily temperatures and validate these predictions using both Monte Carlo simulations and 126 years of available data from the city of Philadelphia. Using extreme statistics, we derive the number and the magnitude of record temperature events, based on the observed Gaussian daily temperature distribution in Philadelphia, as a function of the number of years of observation. We then consider the case of global warming, where the mean temperature systematically increases with time. Over the 126-year time range of observations, we argue that the current warming rate is insufficient to measurably influence the frequency of record temperature events, a conclusion that is supported by numerical simulations and by the Philadelphia data. We also study the role of correlations between temperatures on successive days and find that they do not affect the frequency or magnitude of record temperature events.

PACS numbers: 92.60.Ry, 92.60.Wc, 92.70.-j, 02.50.Cw

I. INTRODUCTION

Almost every summer, there is a heat wave somewhere in the US that garners popular media attention [1]. During such hot spells, daily record high temperatures for various cities are routinely reported in local news reports. A natural question arises: is global warming the cause of such heat waves or are they merely statistical fluctuations? Intuitively, record-breaking temperature events should become less frequent with time if the average temperature is stationary. Thus it is natural to be concerned that global warming is playing a role when there is a proliferation of record-breaking temperature events. In this work, we investigate how systematic climatic changes, such as global warming, affect the magnitude and frequency of record-breaking temperatures. We then assess the potential role of global warming by comparing our predictions both to record temperature data and to Monte Carlo simulation results.

It bears emphasizing that record-breaking temperatures are distinct from threshold events, defined as observations that fall outside a specified threshold of the climatological temperature distribution [2]. Thus, for example, if a city's record temperature for a particular day is 40°C, then an increase in the frequency of daily temperatures above 36°C (*i.e.*, above the 90th percentile) is a threshold event, but not a record-breaking event. Trends in threshold temperature events are also impacted by climate change, and is thus an active research area [2–7]. Studying threshold events is also one of the ways to assess

agricultural, ecological, and human health effects due to climate change [8, 9].

Here we examine the complementary issue of record-breaking temperatures, in part because they are popularized by the media during heat waves and they influence public perception of climate change, and in part because of the fundamental issues associated with record statistics. We focus on daily temperature extremes in the city of Philadelphia, for which data are readily available on the Internet for the period 1874–1999 [10]. In particular, we study how temperature records evolve in time for each *fixed* day of the year. That is, if a record temperature occurs on January 1, 1875, how long until the next record on January 1 occurs? Using the fact that the daily temperature distribution is well approximated by a Gaussian (Sec. II B), we will apply basic ideas from extreme value statistics in Sec. III to predict the magnitude of the temperature jump when a new record is set, as well as the time between successive records on a given day. These predictions are derived for an arbitrary daily temperature distribution, and then we work out specific results for the idealized case of an exponential daily temperature distribution and for the more realistic Gaussian distribution.

Although individual record temperature events are fluctuating quantities, the average size of the temperature jumps between successive records and the frequency of these records are systematic functions of time (see, *e.g.*, [11] for a general discussion). This systematic behavior permits us to make meaningful comparisons between our theoretical predictions, numerical simulations (Sec. IV), and the data for record temperature events in Philadelphia (Sec. V). Clearly, it would be desirable to study long-term temperature data from many locations to discriminate between the expected number of record events for a stationary climate and for global warming. For U.S. cities, however, daily temperature records extend back only 100–140 years [12, 13], and there are both gaps in

*Electronic address: redner@bu.edu; Most of this work was completed when the first author was on leave at the Center for Nonlinear Studies and Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

†Electronic address: mpetersen@lanl.gov

the data and questions about systematic effects caused by “heat islands” for observation points in urban areas. In spite of these practical limitations, the Philadelphia data provide a useful testing ground for our theoretical predictions.

In Sec. VI, we investigate the effect of a slow linear global warming trend [14, 28] on the statistics of record-high and record-low temperature events. We argue that the presently-available 126 years of data in Philadelphia, coupled with the current global warming rate, are insufficient to meaningfully alter the frequency of record temperature events compared to predictions based on a stationary temperature. This conclusion is our main result. Finally, we study the role of correlations in the daily temperatures on the statistics of record temperature events in Sec. VII. Although there are substantial correlations between temperatures on nearby days and record temperature events tend to occur in streaks, these correlations do not affect the frequency of record temperature events for a given day. We summarize and offer some perspectives in Sec. VIII.

II. TEMPERATURE OBSERVATIONS

The temperature data for Philadelphia were obtained from a website of the Earth and Mineral Sciences department at Pennsylvania State University [10]. The data contain both the low and high temperatures in Philadelphia for each day between 1874 and 1999. The data are reported as an integer in degrees Fahrenheit, so we anticipate an error of $\pm 1^\circ\text{F}$. No information is provided about the accuracy of the measurement or the precise location where the temperature is measured. Thus there is no provision for correcting for the heat island effect if the weather station is in an increasingly urbanized location during the observation period. For each day, we also document the middle temperature, defined as the average of the daily high and daily low.

To get a feeling for the nature of the data, we first present basic observations about the average annual temperature and the variation of the temperature during a typical year.

A. Annual averages and extremes

Figure 1 shows the average annual high, middle, and low temperature for each year between 1874 and 1999. To help discern systematic trends, we also plot 10-year averages for each data set. The average high temperature for each year is increasing from 1874 until approximately 1950 and again after 1965, but is decreasing from 1950 to 1965. Over the 126 years of data, a linear fit to the time dependence of the annual high temperature for Philadelphia gives an increase of 1.62°C , compared to the well-documented global warming rate of $0.6 \pm 0.2^\circ\text{C}$ over the past century [9]. On the other hand, there does

not appear to be a systematic trend in the dependence of the annual low temperature on the year. A linear fit to these data give a *decrease* of -0.38°C . This disparity between high and low temperatures is a puzzling and as yet unexplained feature of the data.




FIG. 1: (Color online) Average annual high, middle, and low temperature (in degrees Celsius) for each year between 1874 and 1999 (jagged dotted lines). Also shown are the corresponding 10-year averages (solid curves).

A basic feature about the daily temperature is its approximately sinusoidal annual variation (Fig. 2). The coldest time of the year is early February while the warmest is late July. An amusing curiosity is the discernible small peak during the period January 20–25. This anomaly is the traditional “January thaw” in the northeastern US where sometimes snowpack can melt and a spring-like aura occurs before winter returns (see [15] for a detailed discussion of this phenomenon).




FIG. 2: (Color online) Record high, average high, middle, and low, and record low temperature (in degrees Celsius) for each day of the year.

Also shown, in Fig. 2, are the temperature extremes for each day. The highest recorded temperature in Philadel-

phia of 41.1°C (106°F) occurred on August 7, 1918, while the lowest temperature of -23.9°C (-11°F) occurred on February 9, 1934. Record temperatures also fluctuate more strongly than the mean temperature because there are only 126 years of temperature data. As a result of this short time span, some days of the year have experienced very few records and the resulting current extreme temperature can be far from the value that is expected on statistical grounds (see Sec. V).

B. Daily temperature distribution

To understand the magnitude and frequency of daily record temperatures, we need the underlying temperature distribution for each day of the year. Because temperatures have been recorded for only 126 years, the temperature distribution for each individual day is not smooth. To mitigate this problem, we aggregate the temperatures over a 9-day range and then use these aggregated data to define the temperature distribution for the middle day in this range. Thus, for example, for the temperature distribution on January 5, we aggregate all 126 years of temperatures from January 1–9 (1134 data points). We also use the middle temperature for each day to define the temperature distribution.




FIG. 3: (Color online) Nine-day aggregated temperature distributions for January 5, April 5, October 5, and July 5 in degrees Celsius (top to bottom). Each data set is averaged over a 10% range—10, 9, 8, and 6 points respectively, for January 5, April 5, October 5, and July 5. The distributions are all shifted horizontally by the mean temperature for the day and then vertically to render all curves distinct. The dashed curves are visually-determined Gaussian fits.

Figure 3 shows these aggregated temperature distributions for four representative days—the 5th of January, April, July, and October. Each distribution is shifted vertically to make them all non-overlapping. We also subtracted the mean temperature from each of the dis-

tributions, so that they are all centered about zero. Visually, we obtain good fits to these distributions with the Gaussian $P(\Delta T) \propto e^{-(\Delta T)^2/2\sigma^2}$, where ΔT is the deviation of the temperature from its mean value (in $^\circ\text{C}$), and with $\sigma \approx 5.07, 4.32, 4.12,$ and 3.14 for January 5, April 5, October 5, and July 5, respectively. We therefore use a Gaussian daily temperature distribution as the input to our investigation of the frequency of record temperatures in the next section.

An important caveat needs to be made about the daily temperature distribution. Physically, this distribution cannot be Gaussian *ad infinitum*. Instead, the distribution must cut off more sharply at finite temperature values that reflect basic physical limitations (such as the boiling points of water and nitrogen). We will show in the next section that such a cutoff strongly influences the average waiting time between successive temperature records on a given day.

Notice that the width of the daily temperature distribution is largest in the winter and smallest in the summer. Another intriguing aspect of the daily distributions is the tail behavior. For January 5, there are deviations from a Gaussian at both at the high- and low-temperature extremes, while for April 5 and October 5, there is an enhancement only on the high-temperature side. This enhancement is especially pronounced on April 5, which corresponds to the season where record high temperatures are most likely to occur (see Sec. VIII and Fig. 13). What is not possible to determine with 126 years of data is whether the true temperature distribution is Gaussian up to the cutoff points and the enhancement results from relatively few data, or whether the true temperature distribution on April 5 actually has a slower than Gaussian high-temperature decay.

III. EVOLUTION OF RECORD TEMPERATURES

We now determine theoretically the frequency and magnitude of record temperature events. The schematic evolution of these two characteristics is sketched in Fig. 4 for the case of record high temperatures. Each time a record high for a *fixed* day of the year is set, we document the year t_i when the i^{th} record occurred and the corresponding record high temperature T_i . Under the (unrealistic) assumptions that the temperatures for each day are independent and identical, we now calculate the average values of T_i and t_i and their underlying probability distributions (For a general discussion of record statistics for excursions past a fixed threshold, see *e.g.* [14, 16], while related work on the evolution of records is given in Ref. [17]).

Suppose that the daily temperature distribution is $p(T)$. Two subsidiary distributions needed for record statistics are: (i) the probability that a randomly-drawn temperature *exceeds* T , $p_>(T)$, and (ii) the probability that that this randomly-selected temperature *is less than*




FIG. 4: Schematic evolution of the record high temperature on a specified day for each passing year. Each dot represents the daily high temperature for different years. The first temperature is, by definition, the zeroth record temperature T_0 . This event occurs in year $t_0 = 0$. Successive record temperatures T_1, T_2, T_3, \dots occur in years t_1, t_2, t_3, \dots

$T, p_{<}(T)$. These distributions are [18]:

$$p_{<}(T) \equiv \int_0^T p(T') dT'; \quad p_{>}(T) \equiv \int_T^\infty p(T') dT'. \quad (1)$$

We now determine the k^{th} record temperature T_k recursively. We use the terminology of record high temperatures, but the same formalism applies for record lows. Clearly T_0 coincides with the mean of the daily temperature distribution, $T_0 \equiv \int_0^\infty T p(T) dT$. The next record temperature is the mean value of that portion of the temperature distribution that lies beyond T_0 : that is,

$$T_1 \equiv \frac{\int_{T_0}^\infty T p(T) dT}{\int_{T_0}^\infty p(T) dT}. \quad (2)$$

This formula actually contains a sleight of hand. More properly, we should average the above expression over the probability distribution for T_0 to obtain the true average value of T_1 , rather than merely using the typical or the average value of T_0 in the lower limit of the integral. Equation (2) therefore does not give the true average value of T_1 , but rather gives what we term the *typical* value of T_1 . We will show how to compute the average value shortly.

Proceeding recursively, the relation between successive typical record temperatures is given by

$$T_{k+1} \equiv \frac{\int_{T_k}^\infty T p(T) dT}{\int_{T_k}^\infty p(T) dT}, \quad (3)$$

where the above caveat about using the typical value of T_k in the lower limit, rather than the average over the (as yet) unknown distribution of T_k , still applies.

We now compute $\mathcal{P}_k(T)$, the probability that the k^{th} record temperature equals T ; this distribution is subject to the initial condition $\mathcal{P}_0(T) = p(T)$. For the k^{th} record temperature, the following conditions must be satisfied (refer to Fig. 4): (i) the previous record temperature T' must be less than T , (ii) the next n temperatures, with n arbitrary, must all be less than T' , and (iii) the

last temperature must equal T . Writing the appropriate probabilities for each of these events, we obtain

$$\begin{aligned} \mathcal{P}_k(T) &= \left(\int_0^T \mathcal{P}_{k-1}(T') \sum_{n=0}^{\infty} [p_{<}(T')]^n dT' \right) p(T) \\ &= \left(\int_0^T \frac{\mathcal{P}_{k-1}(T')}{p_{>}(T')} dT' \right) p(T). \end{aligned} \quad (4)$$

The above formula recursively gives the probability distribution for each record temperature in terms of the distribution for the previous record.

Complementary to the magnitude of record temperatures, we determine the time between successive records. Suppose that the current record temperature equals T_k and let $q_n(T_k)$ be the probability that a new record high—the $(k+1)^{\text{st}}$ —is set n years later. For this new record, the first $n-1$ highs after the current record must all be less than T_k , while the n^{th} high temperature must exceed T_k . Thus

$$q_n(T_k) = p_{<}(T_k)^{n-1} p_{>}(T_k). \quad (5)$$

The number of years between the k^{th} record high T_k and the $(k+1)^{\text{st}}$ record T_{k+1} is therefore

$$t_{k+1} - t_k = \sum_{n=1}^{\infty} n p_{<}^{n-1} p_{>} = \frac{1}{p_{>}(T_k)}. \quad (6)$$

We emphasize that this waiting time gives the time between the k^{th} record and the $(k+1)^{\text{st}}$ record when the k^{th} record temperature equals the specified value T_k . If the typical value of T_k is used in Eq. (6), we thus obtain a quantity that we term the typical value of t_k .

To obtain the true average waiting time, we first define $Q_n(k)$ as the probability that the k^{th} record is broken after n additional temperature observations, averaged over the distribution for T_k . Using the definition of q_n , we obtain the formal expression

$$\begin{aligned} Q_n(k) &\equiv \int_0^\infty \mathcal{P}_k(T) q_n(T) dT \\ &= \int_0^\infty \mathcal{P}_k(T) p_{<}(T)^{n-1} p_{>}(T) dT. \end{aligned} \quad (7)$$

Different approaches to determine the Q_n are given in Refs. [14, 19].

There are a number of fundamental results available about record statistics that are *universal* and do not depend on the form of the initial daily temperature distribution, as long as the daily temperatures are independent and identically-distributed (iid) continuous variables [14, 19–22]. In a string of $n+1$ observations (starting at time $n=0$), there are $n!$ permutations of the temperatures out of $(n+1)!$ total possibilities in which the largest temperature is the last of the string. Thus the probability that a new record occurs in the n^{th} year of observation, R_n , is simply [14, 19–22]

$$R_n = \frac{1}{n+1}. \quad (8)$$

In a similar vein, the probability that the initial (0th) record is broken at the n^{th} observation, $Q_n(0)$, requires that the last temperature is the largest while the 0th temperature is the second largest out of $n + 1$ independent variables. The probability for this event is therefore

$$Q_n(0) = \frac{1}{n(n+1)}, \quad (9)$$

again independent of the form of the daily temperature distribution. Thus the average waiting time between the zeroth and first records, $\langle n \rangle = \sum_{n=1}^{\infty} n Q_n(0)$ is infinite!

More generally, the distribution of times between successive records can be obtained by simple reasoning [20, 21]. Consider a string of iid random variables that are labeled by the time index n , with $n = 0, 1, 2, \dots, t$. Define the indicator function

$$\sigma_n = \begin{cases} 1 & \text{if record occurs in } n^{\text{th}} \text{ year} \\ 0 & \text{otherwise.} \end{cases} \quad (10)$$

By definition, the probability for a record to occur in the n^{th} year is $R_n = \langle \sigma_n \rangle = \frac{1}{n+1}$. Therefore the average number of records that have occurred up to time t is

$$\langle R_n \rangle = \sum_{n=1}^t \langle \sigma_n \rangle \sim \ln t. \quad (11)$$

Moreover, because the order of all non-record events is immaterial in the probability for a record event, there are no correlations between the times of two successive record events. That is, $\langle \sigma_m \sigma_n \rangle = \langle \sigma_m \rangle \langle \sigma_n \rangle$. Thus the probability distribution of records is described by a Poisson process in which the mean number of records up to time t is $\ln t$. Consequently, the probability $\Pi(n, t)$ that n records have occurred up to time t is given by [20]

$$\Pi(n, t) \sim \frac{(\ln t)^n}{n!} e^{-\ln t} = \frac{(\ln t)^n}{n!} \frac{1}{t}. \quad (12)$$

To appreciate the implications of these formulae for record statistics, we first consider the warm-up exercise of an exponential daily temperature distribution. For this case, all calculations can be performed explicitly and the results provide intuition into the nature of record temperature statistics. We then turn to the more realistic case of the Gaussian temperature distribution.

A. Exponential distribution

Suppose that the temperature distribution for each day of the year is $p(T) = \mathcal{T}^{-1} e^{-T/\mathcal{T}}$. Equation (1) then gives

$$p_{<}(T) = 1 - e^{-T/\mathcal{T}}; \quad p_{>}(T) = e^{-T/\mathcal{T}}. \quad (13)$$

We now determine the typical value of each T_k . The zeroth record temperature is $T_0 = \int_0^{\infty} T p(T) dT = \mathcal{T}$.

Performing the integrals in Eq. (3) successively for each k gives the basic result

$$T_k = (k+1)\mathcal{T}, \quad (14)$$

namely, a constant jump between typical values of successive record temperatures.

For the probability distribution for each record temperature, we compute $\mathcal{P}_k(T)$ one at a time for $k = 0, 1, 2, \dots$ using Eq. (4). This gives the gamma distribution [23]

$$\mathcal{P}_k(T) = \frac{1}{k!} \frac{T^k}{\mathcal{T}^{k+1}} e^{-T/\mathcal{T}}. \quad (15)$$

This distribution reproduces the typical values of successive temperature records given by Eq. (14); thus the typical and true average values for each record temperature happen to be identical for an exponential temperature distribution. The standard deviation of $\mathcal{P}_k(T)$ is given by $\sqrt{\langle T^2 \rangle - \langle T \rangle^2} = \mathcal{T} \sqrt{k+1}$, so that successive record temperatures become less sharply localized as k increases.

For the typical time between the k^{th} and $(k+1)^{\text{st}}$ records, Eq. (6) gives

$$t_{k+1} - t_k = \frac{1}{p_{>}(T_k)} = e^{T_k/\mathcal{T}}. \quad (16)$$

Substituting $T_k = (k+1)\mathcal{T}$ into Eq. (16), the typical time is $e^{T_k/\mathcal{T}} = e^{(k+1)}$. Thus records become less likely as the years elapse. Notice that the time between records does not depend on \mathcal{T} because of a cancellation between the size of the temperature ‘‘barrier’’ (the current record) and the size of the jump to surmount the record.

For the distribution of waiting times between records, we first consider the time between T_0 and T_1 in detail to illustrate our approach. Substituting Eqs. (13) and (15) into Eq. (7), this distribution is

$$Q_n(0) = \frac{1}{\mathcal{T}} \int_0^{\infty} e^{-T/\mathcal{T}} (1 - e^{-T/\mathcal{T}})^{n-1} e^{-T/\mathcal{T}} dT. \quad (17)$$

Performing this integral by parts gives the result of Eq. (9), $Q_n(0) = 1/[n(n+1)]$.

For later applications, however, we determine the large- n behavior of $Q_n(0)$ by an asymptotic analysis. Defining $x = T/\mathcal{T}$, we rewrite Eq. (17) for large n as

$$\begin{aligned} Q_n(0) &= \int_0^{\infty} e^{-x} (1 - e^{-x})^{n-1} e^{-x} dx \\ &\sim \int_0^{\infty} e^{-2x} e^{-ne^{-x}} dx. \end{aligned} \quad (18)$$

The double exponential in the integrand changes suddenly from 0 to 1 when $n = e^x$, or $x = \ln n$. To estimate $Q_n(0)$, we may omit the double exponential in the integrand and simply replace the lower limit of the integral by $\ln n$. This approach immediately leads to $Q_n(0) \sim n^{-2}$, in agreement with the exact result.

In general, the average waiting time between the k^{th} and $(k+1)^{\text{st}}$ record is, from Eq. (7),

$$Q_n(k) = \int_0^\infty \frac{1}{k!} \frac{T^k}{\mathcal{T}^{k+1}} e^{-T/\mathcal{T}} (1 - e^{-T/\mathcal{T}})^{n-1} e^{-T/\mathcal{T}} dT. \quad (19)$$

While we can express this integral exactly in terms of derivatives of the β function [24], it is more useful to determine its asymptotic behavior by the same analysis as that given in Eq. (18). We thus rewrite $(1 - e^{-x})^{n-1}$ as a double exponential and use the fact that this function is sharply cut off for $x < \ln n$ to reduce the integral of Eq. (19) to

$$Q_n(k) \sim \int_{\ln n}^\infty \frac{x^k}{k!} e^{-2x} dx. \quad (20)$$

To find the asymptotic behavior of this integral, we note that the integrand has a maximum at $x^* = k/2$. Thus for $n > x^*$, the exponential decay term controls the integral and we may again estimate its value by taking the integrand at the lower limit to give $Q_n(k) \propto (\ln n)^k/n^2$. As a result of the power-law tail, the average waiting time between *any* two consecutive records is infinite.

However, the observationally meaningful quantity is the typical value of the waiting time and we thus focus on typical values to characterize the steps between successive records depicted in Fig. 4. The typical time to reach the k^{th} record, t_k , is simply the sum of the typical times between records. Thus

$$\begin{aligned} t_k &= (t_k - t_{k-1}) + (t_{k-1} - t_{k-2}) + \dots + (t_2 - t_1) + t_1 \\ &= e^k + e^{k-1} + \dots + e^2 + e^1 \\ &= \frac{e^k - 1}{1 - e^{-1}} \approx 1.58e^k. \end{aligned} \quad (21)$$

Equivalently, $\ln t_k \approx k + 0.459$ so that Eq. (14) gives $T_k \approx (\ln t_k + 0.541)\mathcal{T}$. Therefore the k^{th} record high temperature increases logarithmically with the total number of observations, as expected from basic extreme statistics considerations [18].

After k record temperatures for a given day have been set, the probability for the next record to occur is $p_>(T_k) = e^{-T_k/\mathcal{T}}$. Since $T_k \approx \mathcal{T} \ln t_k$, we recast this probability as a function of time to obtain

$$p_>(t) = e^{-T_k/\mathcal{T}} \propto e^{-\ln t} = 1/t, \quad (22)$$

thus reproducing the general result in [14, 19–22]. The annual number of record temperatures after t years should be $365/t$; for the Philadelphia data, this gives 2.90 record temperatures for the year 2000, 126 years after the start of observations.

B. Gaussian distribution

We now study record temperature statistics for the more realistic case of a Gaussian daily temperature distribution. Again, to avoid the divergence caused by the unphysical infinite limits in the Gaussian, we begin by computing the typical value T_k of the k^{th} record temperature,

and the typical time t_k until this record. While the calculational steps to obtain these quantities are identical to those of the previous subsection, the details are more complicated because the integrals for $p_<$ and $p_>$ must be evaluated numerically or asymptotically.

As will become evident, the mean value in the Gaussian merely sets the value of T_0 and plays no further role in successive record temperatures. Thus for the daily temperature distribution, we use the canonical form

$$p(T) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-T^2/2\sigma^2} \quad (23)$$

to determine the values of successive record temperatures. The exceedance probability then is

$$\begin{aligned} p_>(T) &= \int_T^\infty \frac{1}{\sqrt{2\pi\sigma^2}} e^{-x^2/2\sigma^2} dx = \frac{1}{2} \operatorname{erfc}\left(T/\sqrt{2\sigma^2}\right) \\ &\sim \frac{1}{\sqrt{2\pi}} \frac{\sigma}{T} e^{-T^2/2\sigma^2} \quad T \gg \sqrt{2\sigma^2}, \end{aligned} \quad (24)$$

where $\operatorname{erfc}(z)$ is the complementary error function [24].

Clearly, $T_0 = 0$, since the Gaussian distribution is symmetric. If we had used a Gaussian with a non-zero mean value, then all the T_k would merely be shifted higher by this mean value. For the next record temperature, Eq. (3) gives

$$T_1 = \frac{\int_0^\infty \frac{1}{\sqrt{2\pi\sigma^2}} T e^{-T^2/2\sigma^2} dT}{\int_0^\infty \frac{1}{\sqrt{2\pi\sigma^2}} e^{-T^2/2\sigma^2} dT}. \quad (25)$$

Substituting $u = T^2/2\sigma^2$ and $v = T/\sqrt{2\sigma^2}$ in the numerator and denominator, respectively, we obtain

$$T_1 = \frac{\int_0^\infty \frac{\sigma}{\sqrt{2\pi}} e^{-u} du}{\frac{1}{2} \operatorname{erfc}(0)} = \sqrt{\frac{2}{\pi}} \sigma. \quad (26)$$

Continuing this recursive computation, Eq. (3) gives

$$\begin{aligned} T_{k+1} &= \frac{\int_{T_k}^\infty \frac{1}{\sqrt{2\pi\sigma^2}} T e^{-T^2/2\sigma^2} dT}{\int_{T_k}^\infty \frac{1}{\sqrt{2\pi\sigma^2}} e^{-T^2/2\sigma^2} dT} \\ &= \frac{T_1 e^{-T_k^2/2\sigma^2}}{\operatorname{erfc}(T_k/\sqrt{2\sigma^2})}. \end{aligned} \quad (27)$$

For the first few k , it is necessary to evaluate the error function numerically and we find $T_2 \approx 1.712T_1$, $T_3 \approx 2.288T_1$, $T_4 = 2.782T_1$, *etc.* Now from Eq. (26), the argument of the error function in Eq. (27) is $T_k/\sqrt{2\sigma^2} = T_k/(T_1\sqrt{\pi})$. Thus for $k \geq 3$, this argument is greater than 1, and it becomes increasingly accurate to use the large- z asymptotic form [24]

$$\operatorname{erfc}(z) \sim \frac{e^{-z^2}}{z\sqrt{\pi}} \left(1 - \frac{1}{2z^2} + \dots\right).$$

This approximation reduces the recursion for T_{k+1} to

$$\begin{aligned} T_{k+1} &= \frac{T_1 e^{-T_k^2/2\sigma^2}}{\operatorname{erfc}(T_k/\sqrt{2\sigma^2})} \\ &\sim \frac{T_1 e^{-T_k^2/2\sigma^2}}{\sqrt{\frac{2\sigma^2}{\pi T_k^2} e^{-T_k^2/2\sigma^2} \left(1 - \frac{1}{2(T_k/\sqrt{2\sigma^2})^2} + \dots\right)}} \\ &\sim T_k \left(1 + \frac{\sigma^2}{T_k^2}\right), \end{aligned} \quad (28)$$

where we have used $T_1 = \sqrt{2\sigma^2/\pi}$ from Eq. (26).

Writing the last line as $T_{k+1} - T_k = \sigma^2/T_k$, approximating the difference by a derivative, and integrating, the k^{th} record temperature for large k has the remarkably simple form

$$T_k \sim \sqrt{2k\sigma^2}. \quad (29)$$

Thus successive record temperatures asymptotically become more closely spaced for the Gaussian distribution. It should be noted, however, that the largest number of record temperature events on any given day in the Philadelphia data is 10, so that the applicability of the asymptotic approximation is necessarily limited.

The more fundamental measure of the temperature jumps is again $\mathcal{P}_k(T)$, the probability distribution that the k^{th} record high equals T . For a Gaussian daily temperature distribution, the general recursion given in Eq. (4) for $\mathcal{P}_k(T)$ is no longer exactly soluble, but we can give an approximate solution that we expect will become more accurate as k is increased. We merely employ the large- T asymptotic form for $p_>(T)$ in the recursion for $\mathcal{P}_k(T)$ even when k is small so that T is not necessarily much larger than σ . Using this approach, we thus obtain for $\mathcal{P}_1(T)$

$$\begin{aligned} \mathcal{P}_1(T) &\approx \left(\int_0^T \frac{\frac{1}{\sqrt{2\pi\sigma^2}} e^{-T'^2/2\sigma^2}}{\sqrt{\frac{\sigma^2}{2\pi T'^2} e^{-T'^2/2\sigma^2}}} dT' \right) \frac{1}{\sqrt{2\pi\sigma^2}} e^{-T^2/2\sigma^2} \\ &\sim \frac{T^2}{2\sigma^2} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-T^2/2\sigma^2}. \end{aligned} \quad (30)$$

Continuing this straightforward recursive procedure then gives

$$\mathcal{P}_k(T) \approx \frac{1}{\Gamma(k + \frac{1}{2})} \frac{T^{2k}}{\sigma^{2k+1}} e^{-T^2/2\sigma^2}, \quad (31)$$

where the amplitude is determined after the fact by demanding that the distribution is normalized.

In spite of the crudeness of this approximation, this distribution agrees reasonably with our numerical simulation results shown in Fig. 5 (details of the simulation are described in the following section). The distributions $\mathcal{P}_k(T)$ move systematically to higher temperatures and become progressively narrower as k increases, in accordance with naive intuition. The approximate form of




FIG. 5: Simulation data for the probability distribution of the k^{th} record high temperature in degrees Celsius, $\mathcal{P}_k(T)$. The distribution $\mathcal{P}_0(T)$ coincides with the Gaussian of Eq. (23), whose parameters match the average temperature and dispersion in Philadelphia. The solid curves correspond to a stationary temperature, while the dashed curves correspond to global warming with rate $v = 0.012^\circ\text{C year}^{-1}$ (see Sec. VI).

Eq. (31) gives a similar shape to the simulated distributions, but there is an overall shift to higher temperatures by roughly $1\text{--}2^\circ\text{C}$.

Next, we study the typical time between successive record temperatures. Equation (6) states that $t_{k+1} - t_k = 1/(p_>(T_k))$. Using the above asymptotic expansion of the complementary error function in the integral for $p_>$ and $T_k \sim \sqrt{2k\sigma^2}$ from Eq. (29), we obtain, for large k ,

$$t_{k+1} - t_k \sim \sqrt{4\pi} \frac{T_k}{\sqrt{2\sigma^2}} e^{T_k^2/2\sigma^2} \sim \sqrt{4\pi k} e^k. \quad (32)$$

Again, the times between records are independent of σ ; this independence arises because both the size of the record and the magnitude of the jumps to surpass the record are proportional to σ , so that its value cancels out in the waiting times.

Finally, we compute the asymptotic behavior for the distribution of waiting times between records. For simplicity, we consider only the waiting time distribution $Q_n(0)$ until the first record. The distribution of waiting times for subsequent records has the same asymptotic tail as $Q_n(0)$, but also contains more complicated pre-asymptotic factors. Substituting the Gaussian for $p(T)$ and the asymptotic form for $p_>(T)$ into Eq. (7), and then expanding $(1-p_>)^{n-1}$ as a double exponential, we obtain

$$Q_n(0) \sim \int_0^\infty \frac{1}{2\pi x} \exp\left[-\frac{x^2}{\sigma^2} - \sqrt{\frac{n^2\sigma^2}{2\pi x^2}} e^{-x^2/2\sigma^2}\right] dx. \quad (33)$$

The double exponential again cuts off the integral when x is less than a threshold value $x^* \sim \sqrt{2\sigma^2 \ln n}$. As a

result, Eq. (33) reduces to

$$Q_n(0) \sim \int_{\sqrt{2\sigma^2 \ln n}}^{\infty} \frac{1}{2\pi x} e^{-x^2/\sigma^2} dx \sim \frac{1}{n^2}. \quad (34)$$

In the final result, we drop logarithmic corrections because the approximation made in writing Eq. (33) also contains errors of the same magnitude. Thus the distribution of waiting times n until the first record again has a n^{-2} power-law tail and the mean waiting time is infinite.

The typical time until the k^{th} record is again given by the sum of successive time intervals. Asymptotically, Eq. (32) gives

$$t_k \sim \int_0^k \sqrt{4\pi n} e^n dn \sim \sqrt{4\pi k} e^k, \quad (35)$$

or $k \approx \ln t - \frac{1}{2} \ln(4\pi \ln t)$. Thus the number of records grows slowly with time; this result has the obvious consequence that records become less likely to occur at later times.

IV. MONTE CARLO SIMULATIONS

To verify our theoretical derivations, Monte Carlo simulations were performed for both the exponential and Gaussian temperature distributions. Our simulations typically involve 10^5 realizations (days) over a minimum of 1000 years of observations, and continue until six record temperatures have been achieved. We use “years” consisting of 10^5 days so that we generate a sufficient number of record temperatures to have reasonable statistics. For our initial simulations, we used a stationary mean and variance of 18°C and 5°C respectively, which are typical values for the distribution of maximum daily temperatures in the spring or fall in Philadelphia. However, the numerical validation of our theoretical distributions does not depend on the particular values of mean and variance.

The simulation errors using an exponential distribution for the k^{th} record (with $k = 0, 1, \dots, 5$) are: less than 3×10^{-5} for $\mathcal{P}_k(T)$ (Eq. (15)) using a distribution with 100 bins; 8.3×10^{-5} for $Q_n(0)$ (Eq. (9)); 2.2×10^{-3} (relative error) for the mean temperature of the k^{th} record temperature (Eq. (14)); and 0.01 (relative error) for the variance. The Gaussian distribution yields fewer exact expressions for comparison, but includes a relative error of 6.4×10^{-3} for the mean temperature of the k^{th} record temperature (Eq. (28)), $k = 0 \dots 5$. For both the exponential and Gaussian distributions, the probability of breaking a record temperature with time is well fit by the form $1/(t+1)$, with an error of less than 9.2×10^{-5} . These errors decrease as the number of realizations increases, and the small errors for simulations with 10^5 realizations confirm the correctness of the theoretical distributions.

Monte Carlo simulations were also performed to explore the effect of temporal correlations in daily temperatures on the frequency statistics of record-temperature

events and the magnitude of successive record temperatures. This topic will be discussed in detail in Sec. VII. We used the Fourier filtering analysis method [25, 26] to generate power-law correlations between daily temperature data for years consisting of 10^4 days over 200 years and for several values of the exponent in the power law of the temporal correlation function.

V. RECORD TEMPERATURE DATA

Between 1874 and 1999, a total of 1707 record highs (4.68 for each day on average) and 1343 record lows (3.68 for each day) occurred in Philadelphia [27]. Because the temperature was reported as an integer, a temperature equaling a current record could represent a new record if the measurement was more accurate. With the less stringent definition that a new record either exceeds or *equals* the current record, the number of record high and record low events over 126 years increased from 1707 to 2126 and from 1343 to 1793, respectively. However, this alternative definition does not qualitatively change the statistical properties of record temperature events.




FIG. 6: (Color online) Average k^{th} record high (Δ) and record low (∇) temperature for each day, divided by the daily temperature dispersion, versus k (from the Philadelphia temperature data). The dashed curve is $T_k/\sigma = 1.15\sqrt{k}$.

To compare with our theory, first consider the size of successive record temperatures. According to Eq. (29), the k^{th} record high (and record low) temperature should be proportional to $\sqrt{2k\sigma^2}$. Because the mean temperature for each day has already been subtracted off, here T_k denotes the absolute value of the difference between the k^{th} record temperature and the zeroth record. To have a statistically meaningful quantity, we compute T_k/σ_α for each day of the year, and then average over the entire year; here the subscript $\alpha = h, l$ denotes the daily dispersion for the high and low temperatures, respectively. As




FIG. 7: Probability that a record high temperature (top) or record low (bottom) occurs at a time t (in years) after the start of observations. The symbols \triangle and ∇ are 10-point averages of Philadelphia data from 1874 to 1999 for ease of visualization. Simulated data were produced by a stationary Gaussian distribution ($v = 0$), or where the mean increases according to $v = 0.003, 0.006$, or 0.012C year^{-1} . The stationary data fit the theoretical expectation of $1/(t+1)$ (thick dashed line), while warming leads the distribution to asymptote to a constant probability (thin dashed lines).

shown in Fig. 6, the annual average for T_k/σ_α is consistent with \sqrt{k} growth for both the record high and record low temperature. Up to the 6th record, both data sets are quite close, and where the data begin to diverge, the number of days with more than 6 records is small—69 for high temperatures and 26 for low temperatures.

Finally, we study the evolution of the frequency of record temperature days as a function of time. As discussed in Sec. III, the number of records in the t^{th} year of observation (since 1874) should be $365/t$. In spite of the year-to-year fluctuations in the number of records, the prediction $365/t$ fits the overall trend (Fig. 7). We also examine the distribution of waiting times between records. Since the amount of data is small, it is useful to study the cumulative distribution, $\mathcal{Q}_n(k) \equiv \sum_{m=n}^{\infty} Q_m(k)$, defined as the probability that




FIG. 8: Probability that the k^{th} record high temperature occurs at time t (in years) or later, using simulated data (solid curves). The $k = 1$ simulated data closely match the asymptotic theoretical distribution of $1/t$ (dashed line). Also shown are the $k = 1$ data for record high temperatures (\triangle) and record low temperatures (∇) for the Philadelphia data.

the time between the k^{th} and the $(k+1)^{\text{st}}$ record temperatures on a given day is n years or larger. As shown in Fig. 8, the agreement between the Philadelphia data and the theoretical prediction from Eq. (34), $\mathcal{Q}_n(0) \propto 1/n$ is quite good. The Monte Carlo simulations match the theoretical prediction nearly exactly, with an rms error of 9×10^{-5} .

In summary, the data for the magnitude of temperature jumps at each successive record, the frequency of record events, and the distribution of times between records are consistent with the theoretical predictions that arise from a Gaussian daily temperature distribution with a stationary mean temperature.

VI. SYSTEMATICALLY CHANGING TEMPERATURE

We now study how a systematically changing average temperature affects the evolution of record temperature events. For global warming, we assume that the mean temperature has a slow superimposed time dependence vt , with $v > 0$ and where t is the time (in years) after the initial observational year.

A. Exponential distribution

Again, as a warm-up exercise, we first consider the idealized case of an exponential daily temperature distribution,

$$p(T; t) = \begin{cases} e^{-(T-vt)} & T > vt \\ 0 & T < vt, \end{cases} \quad (36)$$

where we set the characteristic temperature scale \mathcal{T} to 1 for simplicity. In these units, both T and vt are dimensionless. With this distribution, the recursion Eq. (3) for successive record temperatures becomes

$$T_{k+1} \equiv \frac{\int_{T_k}^{\infty} T e^{-(T-vt_{k+1})} dT}{\int_{T_k}^{\infty} e^{-(T-vt_{k+1})} dT}. \quad (37)$$

The factor $e^{vt_{k+1}}$ appears in both the numerator and denominator and thus cancels. As a result, $T_k = k+1$, *independent* of v . Thus a systematic temperature variation—either global warming or global cooling—does not affect the magnitude of the jumps in successive record high temperatures. This fact was verified by numerical simulations with an exponential distribution, where the distributions of $\mathcal{P}_k(T)$ for $v = 0.012^\circ\text{C year}^{-1}$ and $v = 0$ match to within a few percent for $k = 0 \dots 5$.

On the other hand, a systematic temperature dependence does affect the time between records. Suppose that the current record high temperature of T_k was set in year t_k . Then the exceedance probability at time $t_k + j$ is

$$\begin{aligned} p_{>}(T_k; t_k + j) &= \int_{T_k}^{\infty} e^{-[T-v(t_k+j)]} dT \\ &= e^{-(T_k-vt_k)} e^{jv} \equiv X e^{jv}. \end{aligned} \quad (38)$$

The exceedance probability is thus either enhanced or suppressed by a factor e^v due to global warming or cooling, respectively, for each elapsed year. The probability $q_n(T_k)$ that a new record high temperature occurs n years after the previous record T_k at time t_k is

$$q_n(T_k) = e^{nv} X \prod_{j=1}^{n-1} (1 - e^{jv} X), \quad (39)$$

with $q_1(T_k) = e^v X$; this generalizes Eq. (5) to incorporate a global climatic change.

For the case of global warming ($v > 0$), each successive term in the product decreases in magnitude and there is a value of j for which the factor $(1 - e^{jv} X)$ is no longer positive. At this point, the next temperature must be a new record. Thus we (over)estimate the time until the next record after T_k by the criterion $(1 - e^{jv} X) = 0$, or $j = (T_k - vt_k)/v \sim (k/v) - t_k$. Since this value of j also coincides with $t_{k+1} - t_k$ by construction, we obtain $t_k \sim k/v$. Thus the time between consecutive records asymptotically varies as $t_{k+1} - t_k \sim 1/v$. This conclusion agrees with a previous mathematical proof of the constancy of the rate of new records when a linear temporal trend is superimposed on a set of continuous iid variables [28]; a different approach to deal with a linear trend is given in [29].

If global warming is slow, the waiting time between records will initially increase exponentially with k , as in the case of a stationary temperature, but then there will be a crossover to the asymptotic regime where the waiting time is constant. We estimate the crossover time by equating the two forms for the waiting times, $t_{k+1} - t_k =$

$e^{(k+1)}$ (stationary temperature) and $t_{k+1} - t_k = 1/v$ (increasing temperature), to give $k^* \approx -\ln v$. Now the average annual high temperature in Philadelphia has increased by approximately 1.94°C over 126 years. The resulting warming rate of $0.0154^\circ\text{C per year}$ then gives $k^* \approx 3.6$. Thus the statistics of the first 3.6 record high temperatures should be indistinguishable from those in a stationary climate, after which record temperatures should occur at a constant rate. Since the average number of record high temperatures for a given day is 4.7 and the time until the next record high is very roughly $e^{5.7} - e^{4.7} \approx 190$ years, we are still far from the point where global warming could have an unambiguous effect on the frequency of record high temperatures.

For global cooling ($v < 0$), the waiting time probability becomes

$$q_n(T_k) = \prod_{j=1}^{n-1} (1 - e^{-jw} Y) e^{-nw} Y, \quad (40)$$

with $q_1(T_k) = e^{-w} Y$, where $w \equiv |v|$ is positive, and $Y = e^{-T_k - wt_k}$. We estimate the above product by the following simple approach. When $jw < 1$, then $e^{-jw} \ll 1$, and each factor within the product is approximately $(1 - Y)$. Consequently, for $nw > 1$, each term in the product approximately equals $(1 - Y)$ for $j < n^* = 1/w$, while for $j > n^*$, $e^{-jw} \approx 0$, and the later terms in the product are all equal to 1. Thus

$$q_n(T_k) \sim \begin{cases} (1 - Y)^n e^{-nw} Y & n < n^* \\ (1 - Y)^{1/w} e^{-nw} Y & n > n^* \end{cases}. \quad (41)$$

Using this form for q_n , we find, after straightforward but slightly tedious algebra, that the dominant contribution to the waiting time until the next record temperature, $t_{k+1} - t_k = \sum_{n=1}^{\infty} n q_n$, comes from the terms with $n < n^*$ in the sum. For the case slow global cooling, we thereby find

$$\begin{aligned} t_{k+1} - t_k &\sim \frac{1/Y}{[1 + w(1/Y - 1)]^2} \sim 1/Y \\ &= e^{T_k + wt_k}. \end{aligned} \quad (42)$$

Since $t_{k+1} - t_k \approx dt/dk$ and using $T_k \sim k$, Eq. (42) can be integrated to give $(1 - e^{-wt_k}) = w(e^k - 1)$. As long as the right-hand side is less than 1, a solution for t_k exists. In the converse case, there is no solution and thus no additional record highs under global cooling, or equivalently, no more record lows for global warming. For small w and in the pre-crossover regime where $e^k \approx t_k$, the criterion for no more records reduces to $t > 1/w$. If the daily low temperature in Philadelphia also experienced a warming rate of $0.0154^\circ\text{C per year}$, then there should be no additional record low temperatures after about 36 years of observations. However, the daily low temperatures do not show a long-term systematic variation, so new record lows should continue to occur, as is observed.

B. Gaussian distribution

We now treat the more realistic case where a systematic temperature variation is superimposed on a Gaussian daily temperature distribution, as embodied by

$$p(T; t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(T-vt)^2/2\sigma^2}. \quad (43)$$

The details of the effects of a systematic temperature variation on the statistics of record temperatures are tedious and we merely summarize the main results. We assume a slow systematic variation, $T_k - vt \gg 0$, so that an asymptotic analysis will be valid. Under this approximation, both global warming or global cooling lead to the following recursion for T_k , to leading order:

$$T_{k+1} - T_k \sim \frac{\sigma^2}{T_k} \left(1 + \frac{vt}{T_k} \right). \quad (44)$$

The term proportional to vt in Eq. (44) is subdominant, so that T_k still scales as $\sim \sqrt{2k\sigma^2}$, for both global warming and global cooling.

Next we determine the times between successive record high temperatures. The basic quantity that underlies these waiting times is again the exceedance probability, when the current record is T_k and the current time is $t_k + j$. Following Eq. (24), this exceedance probability is

$$p_{>}(T_k; t_k + j) \sim \frac{1}{2} \operatorname{erfc} \left(\frac{T_k - v(t_k + j)}{\sqrt{2\sigma^2}} \right). \quad (45)$$

In the asymptotic limit where the argument of the complementary error function is large, the controlling factor in $p_{>}$ is

$$e^{-[T-v(T_k+j)]^2/2\sigma^2} \approx e^{-(T-vt_k)^2/2\sigma^2} e^{vj(T-vt_k)/\sigma^2}. \quad (46)$$

The crucial point is that the latter form for the exceedance probability has the same j dependence as in the exponential distribution (Eq. (38)). Thus our arguments for the role of global warming with an exponential daily temperature distribution continue to apply. In particular, the time between successive records initially grows as $\sqrt{4\pi k} e^k$, but then asymptotically approaches the constant value $1/v$. As a result, the time before global warming measurably influences the frequency of record high and record low temperatures will be similar for both the exponential and Gaussian temperature distributions.

Monte Carlo simulations were performed for warming rates $v = 0.003, 0.006, \text{ and } 0.012^\circ\text{C}/\text{year}$, where the middle case corresponds to the accepted rate of global mean warming of 0.6°C for the 20th century [9]. Unlike the exponential distribution simulations, for the Gaussian distribution $\mathcal{P}_k(T)$ is slightly different in the cases of no warming and warming (Fig. 5).

Figure 7 shows the results of numerical simulations using the Gaussian distribution with 10^5 realizations for the

three warming rates. For the stationary case ($v = 0$), the probability of breaking a record after t years closely follows the theoretical expectation of $1/(t + 1)$. For warming, the rate of breaking a record high (Fig. 7, top) ultimately asymptotes to a constant frequency of approximately $1.25v$ by 10^4 years. Given our crude calculation following Eq. (39) that the time between records is $1/v$, the agreement between the observed rate of $1.25v$ and our estimate of v is gratifying. As also predicted in our theory, the probability of breaking a record low temperature under global warming precipitously decays after a few hundred years (Fig. 7, bottom); eventually record low temperatures simply stop occurring in a warming world.

VII. ROLE OF TEMPORAL CORRELATIONS

Thus far our presentation has been based on *independent* daily temperatures—no correlations between temperatures on successive days. However, from common experience we know that local weather consists of multi-day patterns within which smaller temperature variations occur. Anecdotally, the temperature tomorrow will be close to the temperature today. In fact, it has been found in global climatological data that correlations between temperatures on two widely separated days decay as a power law in the separation [30]. Here we quantify these correlations for the Philadelphia data and then discuss the potential ramifications of these correlations on the frequency of record temperature events.

A. Daily temperature correlation data

From the Philadelphia data, we compute the normalized interday temperature correlation function defined as

$$c_\alpha(i, j) = \frac{\langle T_i T_j \rangle - \langle T_i \rangle \langle T_j \rangle}{\langle T_i^2 \rangle - \langle T_i \rangle^2}. \quad (47)$$

Here i and $j > i$ denote the i^{th} and j^{th} days of the year, T_i is the temperature on the i^{th} day, and $\langle T_i \rangle$ is its average value over the 126 years of data, while the index $\alpha = h, m, l$ denotes the high, middle, and low temperature for each day. If i is a day near the end of the year, then T_j will refer to a temperature in the following year when the separation between the two days exceeds $(365 - i)$. According to Eq. (47), if the temperatures T_i and T_j are both greater than or both less than the respective average temperatures for days i and j , then there is a positive contribution to the correlation function. Thus $c_\alpha(i, j)$ measures systematic temperature deviations from the mean on these two days. For convenience, we normalize the c_α so that they all equal 1 when $|i - j| = 0$.

The correlation functions depend primarily on the separation between the two days, $|i - j|$, and weakly on the initial day i . To obtain a succinct measure of the temperature correlation over a year, we define the annual




FIG. 9: (Color online) The correlation functions $C_\alpha(t)$ for high (Δ), middle (\circ), and low temperature (∇) versus time (in years). The straight line of slope $-4/3$ is a guide for the eye.

average correlation function

$$C_\alpha(t) \equiv \sum_{i=1}^{365} c_\alpha(i, i+t). \quad (48)$$

All three correlations functions are consistent with a power law decay $C_\alpha(t) \sim t^{-\gamma}$ (Fig. 9). Over a range of approximately 1–20 days, the best fit value of γ is 1.29 for C_h (which remains strictly positive until 36 days) and $\gamma = 1.44$ for C_m (which remains strictly positive until 41 days). The correlation function C_l is visibly distinct and remains strictly positive until 149 days, with a best-fit exponent of $\gamma = 1.36$. These power-law decays in the temperature correlation functions are consistent with the previous results of Ref. [30]. However, the exponent value that we observe, approximately $4/3$, is considerably larger than that reported in Ref. [30]. The time integrals of the high-, middle-, and low-temperature correlation functions are 1.78, 2.04 and 5.16 respectively. We may therefore view 1.78 as the average length of an independent high-temperature event, and correspondingly $365/1.78 \approx 205$ as the number of effective independent “days” for high temperatures. Parallel results hold for middle and low temperatures. These numbers provide a feeling for the extent of multiday weather patterns because of temperature correlations.

B. Simulations with correlated daily temperatures

To determine if these correlations affect the frequency and magnitude of record temperature events, we performed Monte Carlo simulations in which daily temperatures had temporal correlations that matched the data discussed above. We generate such correlated data using the Fourier filtering method of Refs. [25, 26] with a

correlation function of the form

$$C_\alpha(t) = t^{-\gamma} \quad (49)$$

for a range of γ values around the observed value of 1.3–1.4. Due to the computational demands of generating correlated data, simulations of years consisting of 10^4 days for 200 years were performed, which are less extensive than our simulations for uncorrelated temperatures. We find that the statistics of the time between record temperature events and the magnitude of successive record temperatures are virtually identical to those obtained when the temperature is an independent identically-distributed random variable. (Figs. 10 and 11). Our results are also not sensitive to the value of the decay exponent γ of the correlation function, within our tested range of $\gamma \in [0.5, 1.5]$. We conclude that the discussion in Secs. III–VI, which assumed uncorrelated day-to-day temperatures, can be applied to real atmospheric observations, where daily temperatures are correlated. It is worth mentioning, however, that interday correlations do strongly affect the statistics of successive extremes in temperatures [31].




FIG. 10: Simulation data for the probability distribution of the k^{th} record high temperature in degrees Celsius, $\mathcal{P}_k(T)$, where daily temperatures are uncorrelated (solid line) and power-law correlated with exponent 1.5. (dashed line).

C. Correlations between record temperature events

While temperature correlations do not affect record statistics for a given day, these correlations should cause records to occur as part of a heat wave or a cold snap, rather than being singular one-day events. As a matter of curiosity, we studied the distribution of times (in days) between successive record events, as well as the distribution of streaks (consecutive days) of record temperatures from the time history of all record temperature events.

Because the number of record temperatures decreases from year to year, these time and streak distributions are




FIG. 11: Probability that the k^{th} record high temperature occurs at time t (in years) or later, using uncorrelated (solid line) and power-law correlated daily temperatures (dashed line).

not stationary. We compensate for this non-stationarity by rescaling so that data for all years can be treated on the same footing. For example, for the distribution of times between successive records, we rescale each interevent time by the average time between records for that year. Thus, for example, if two successive records occurred 78 days apart in a year where 5 record temperature events occurred (average separation of 73 days), the scaled separation between these two events is $\tau = 78/73 \approx 1.068$. For the length of record streaks, we similarly rescale each streak by the average streak length in that year, assuming record temperature events were uncorrelated.




FIG. 12: (Color online) Distribution of times $p(\tau)$ between successive record temperature events (\triangle record highs, ∇ record lows). The times are scaled by the average time between record events for each year.

The distribution of times between successive record temperature days decays slower than exponentially (Fig. 12); the latter form would occur if record temper-

ature events were uncorrelated. In a similar vein, we observe an enhanced probability for records to occur in streaks. Since record streaks are rare, we can only make the qualitative statement that the streak distribution is different than that from uncorrelated data. Our basic conclusion is that interday temperature correlations do affect statistical features of successive record temperature events but do not affect the statistics of record temperatures on a given day, where events are more than one year apart.

VIII. DISCUSSION

Two basic aspects of record temperature events are the size of the temperature jump when a new record occurs and the separation in years between successive records on a given day. We computed the distribution functions for these two properties by extreme statistics reasoning. For the Gaussian daily temperature distribution, we found that (i) the k^{th} record high temperature asymptotically grows as $\sqrt{k} \sigma$, where σ is the dispersion in the daily temperature, and (ii) record events become progressively less likely, with the typical time between the k^{th} and $(k+1)^{\text{st}}$ record growing as $\sqrt{k} e^k$. This latter result is independent of σ so that systematic changes in temperature variability should not affect the time between temperature records.

From these predictions, the distribution of waiting times between two successive records on a given day has an inverse-square power-law tail, with a divergent average waiting time. Furthermore, the number of record events in the t^{th} year of observations decays as t^{-1} [14, 19–22]. These theoretical predictions agree with numerical simulations and with data from 126 years of observations in Philadelphia. Another important feature is that the annual frequency of record temperature events is not measurably influenced by interday power-law temperature. However, these correlations do play a significant role at shorter time scales.

Our primary result is that we cannot *yet* distinguish between the effects of random fluctuations and long-term systematic trends on the frequency of record-breaking temperatures with 126 years of data. For example, in the 100th year of observation, there should be $365/100 = 3.65$ record-high temperature events in a stationary climate, while our simulations give 4.74 such events in a climate that is warming at a rate of 0.6°C per 100 years. However, the variation from year to year in the frequency of record events after 100 years is larger than the difference of $4.74 - 3.65$, which should be expected because of global warming (Fig. 7). After 200 years, this random variation in the frequency of record events is still larger than the effect of global warming. On the other hand, global warming already does affect the frequency of extreme temperature events that are defined by exceeding a fixed threshold [2–7].

While the agreement between our theory and the data for record temperature statistics is satisfying, there are




FIG. 13: (Color online) Number of high-temperature records for each day of the year, averaged over a 30-day range (top). Below are the variances in the high (Δ), middle (\circ), and low (∇) temperatures for each day averaged over a 10-day range.

various facts that we have either glossed over or ignored. These include: (i) a significant difference between the number of record high and record low events—1705 record high events and only 1346 record low events have occurred the 126 years of data. (ii) A propensity for record high temperatures in the early spring. This seasonality is illustrated both by the number of records for each day of the year and by the daily temperature variance $\sigma_i \equiv \sqrt{\langle T_i^2 \rangle - \langle T_i \rangle^2}$, where $\langle T_i \rangle$ and $\langle T_i^2 \rangle$ are the mean and mean-square temperatures for the i^{th} day (Fig. 13). (iii) The potential role of a systematically increasing variability on the frequency of records. For the last point, Krug [32] has shown that for an exponential daily temperature distribution whose width is increasing linearly with time, the number of record events after t years grows as $(\ln t)^2$, intermediate to the $\ln t$ growth of a stationary distribution and linear growth when the aver-

age temperature systematically increases. (iv) Day/night or high/low asymmetry [33]. That is, as a function of time there are more days whose highs exceeds a given threshold and fewer days whose high is less than a threshold. Paradoxically, however, there are fewer days whose lows exceed a given temperature and more days whose lows are less than a given temperature. Since highs generally occur in daytime and lows in nighttime, these results can be restated as follows: the number of hot days is increasing *and* the number of cold nights is increasing. We don't know how this latter statement fits with the phenomenon of global warming.

Another caveat is that our theory applies in the asymptotic limit, where each day has experienced a large number of record temperatures over the observational history. The fact that there are no more than 10 record events on any single day means that we are far from the regime where the asymptotic limit truly applies. Finally, and very importantly, it would be useful to obtain long-term temperature data from many stations to provide a more definitive test of our predictions.

Acknowledgments

We thank D. ben-Avraham for insightful discussions, C. Forest for helpful advice, and P. Huybers and R. Katz for constructive manuscript suggestions and literature advice. We also thank J. Krug for informing us of Refs. [28, 29] about the effect of linear trends on record statistics after this work was completed, S. Majumdar for helpful discussions that led to the derivations given in Eqs. (10)–(12), Z. Racz for making us aware of literature on high/low asymmetry, and a referee for constructive suggestions. Finally, we gratefully acknowledge financial support from DOE grant W-7405-ENG-36 (at LANL) and NSF grants DMR0227670 and DMR0535503 (at BU).

-
- [1] Record-Breaking Heat: Is Global Warming to Blame? J. Roach, National Geographic News, August 16, 2005; http://news.nationalgeographic.com/news/2005/08/0816_050816_heat_wave.html.
- [2] Z. Yan, P. D. Jones, T. D. Davies, A. Moberg and H. Bergstroem, *Climatic Change* **53**, 355 (2002).
- [3] L. O. Mearns, R. W. Katz, and S. H. Schneider, *J. Climate Appl. Meteor.* **23**, 1601 (1984).
- [4] J. Hansen, I. Fung, A. Lacis, D. Rind, S. Lebedeff, R. Ruedy, G. Russel, and P. Stone, *J. Geophys. Res.* **93**, 9341 (1988).
- [5] R. Katz and B. Brown, *Climatic Change* **21**, 289 (1992).
- [6] A. F. Colombo, D. Etkin and B. W. Karney, *J. Climate*, **12**, 2490 (1999).
- [7] M. Unkašević, D. Vujović, and I. Tošić, *Theor. Appl. Climatol.* **82**, 199 (2005).
- [8] G. Meehl, T. Karl, D. Easterling and S. Changnon, *Bull. Am. Met. Soc.* **81**, 413 (2000).
- [9] For a comprehensive discussion of the global warming of $0.6 \pm 0.2^\circ\text{C}$ in the past century, see *e.g.*, IPCC, 2001: *Climate Change 2001: The Scientific Basis*. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change eds. J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Dai, K. Maskell, and C. A. Johnson, (Cambridge University Press, Cambridge, 2001).
- [10] http://www.ems.psu.edu/PA_Climatologist/philadelphia/phlcomp.html.
- [11] For a general reference on the use of statistical methods in climatology, see *e.g.*, H. von Storch and F. W. Zwiers, *Statistical Analysis in Climate Research*, (Cambridge University Press, Cambridge, UK, 1999).

- [12] C. H. Reitan and J. M. Moran, *Monthly Weather Review*, **105** 1442, (1977).
- [13] R. C. Balling, Jr., J. A. Skindlov, and D. H. Phillips, *J. Climate* **3**, 1491 (1990).
- [14] B. C. Arnold, N. Balakrishnan, and H. N. Nagaraja, *Records* (John Wiley & Sons, Inc., New York, 1998).
- [15] C. M. Godfrey, D. S. Wilks, and D. M. Schultz, *Bull. Amer. Meteor. Soc.* **83**, 53 (2002).
- [16] E. Vanmarcke, *Random Fields: Analysis and Synthesis*, (MIT Press, Cambridge, MA, 1983). The original reference is S. O. Rice, *Bell Syst. Tech. J.*, **23**, 282 (1944); **24**, 46 (1945).
- [17] For a related pedagogical analysis of record temperatures, see B. Schmittmann and R. K. Zia, *Am. J. Phys.* **67**, 1269 (1999).
- [18] See *e.g.*, E. J. Gumbel, *Statistics of Extremes* (Dover Publications, Mineola, N.Y., 2004); J. Galambos, *The Asymptotic Theory of Extreme Order Statistics* (Krieger Publishing Co., Malabar, FL, 1987) for a general introduction to extreme statistics.
- [19] N. Glick, *Am. Math. Monthly*, **85**, 2 (1978).
- [20] P. Sibani and P. Littlewood, *Phys. Rev. Lett.* **71**, 1482 (1993); P. Sibani, M. Brandt, and P. Alstrom, *Int. J. Mod. Phys. B* **12**, 361 (1998).
- [21] K. Jain and J. Krug, *J. Stat. Mech.* 04008 (2005); J. Krug and K. Jain, *Physica A* **358**, 1 (2005).
- [22] R. E. Benestad, *Climate Res.* **25**, 3 (2003).
- [23] See <http://mathworld.wolfram.com/GammaDistribution.html>
- [24] M. Abramowitz, and I. A. Stegun *Handbook of Mathematical Functions* (Dover, New York, 1972).
- [25] D. ben-Avraham and S. Havlin *Diffusion and Reactions in Fractals and Disordered Systems*, (Cambridge University Press, Cambridge, England, 2000), Appendix D4.
- [26] H. A. Makse, S. Havlin, M. Schwartz, and H. E. Stanley, *Phys. Rev. E* **53**, 5445 (1996).
- [27] S. Redner and M. Petersen, unpublished. All supplemental data are freely available on request.
- [28] R. Ballerini and S. I. Resnick, *J. Appl. Prob.* **22**, 487 (1985); *Adv. Appl. Prob.* **19**, 801 (1987); K. Borovkov, *J. Appl. Prob.* **36**, 668 (1999). See also Sec. 6.4 in Ref. [14].
- [29] G. W. Bassett, *Climatic Change* **21**, 303 (1992).
- [30] E. Koscielny-Bunde, A. Bunde, S. Havlin, H. E. Roman, Y. Goldreich, and H.-J. Schellnhuber *Phys. Rev. Lett.* **81**, 729 (1998); J. F. Eichner, E. Koscielny-Bunde, A. Bunde, S. Havlin, and H.-J. Schellnhuber, *Phys. Rev. E* **68**, 046133 (2003); A. Bunde, J. F. Eichner, J. W. Kantelhardt, and S. Havlin, *Phys. Rev. Lett.* **94**, 048701 (2005).
- [31] J. F. Eichner, J. W. Kantelhardt, A. Bunde, and S. Havlin, *Phys. Rev. E* **73**, 016130 (2006).
- [32] J. Krug, private communication.
- [33] M. Pattantyús-Ábrahám, A. Király, and I. M. Jánosi, *Phys. Rev. E* **69**, 021110 (2004); D. R. Easterling et al. *Science* **277**, 364 (1997).